

Ethical considerations

The authors declare that the present article contains no personal information that could identify the patient, because it is a review of a clinical case record, no authorization by an ethics committee was needed.

Financial disclosure

No financial support was received in relation to this article.

Conflict of interest

The authors declare that there is no conflict of interest.

References

1. Sigala-Robles R, Aguayo-Patrón SV, Calderón-de la Barca AM. Genética, ambiente y asma asociados a enfermedad celíaca en la familia extendida de un niño afectado. *Rev Gastroenterol Méx.* 2018;83:79–85, <http://dx.doi.org/10.1016/j.rgmx.2017.05.006>.
2. Remes-Troche JM, Núñez-Alvares C, Uscanga-Domínguez LF. Celiac disease in Mexican population: an update. *Am J Gastroenterol.* 2013;108:283–4, <http://dx.doi.org/10.1038/ajg.2012.408>.
3. Hujoel IA, Reilly NR, Rubio Tapia A. Celiac disease: clinical features and diagnosis. *Gastroenterol Clin North Am.* 2019;48:19–37, <http://dx.doi.org/10.1016/j.gtc.2018.09.001>.
4. Cakir M, Cezaroglu S, Cobanoglu U. Celiac disease in children with chronic constipation. *Turk J Med Sci.* 2016;46:651–6, <http://dx.doi.org/10.3906/sag-1502-130>.
5. Husby S, Koletzko S, Korponay-Szabó I, et al. European Society for pediatric gastroenterology hepatology and nutrition guidelines for the diagnosis of coeliac disease 2020. *J Pediatr Gastroenterol Nutr.* 2020;70:141–56, <http://dx.doi.org/10.1097/MPG.0000000000002497>.
6. Remes-Troche JM, Uscanga-Domínguez LF, Aceves-Tavares RG, et al. Guía clínica para diagnóstico y tratamiento de la enfermedad celíaca en México. *Rev Gastroenterol Méx.* 2018;83:434–50, <http://dx.doi.org/10.1016/j.rgmx.2018.05.005>.

7. Ierardi E, Losurdo G, Piscitelli D, et al. Seronegative celiac disease: where is the specific setting? *Gastroenterol Hepatol Bed Bench.* 2015;8:110–6.
8. Singh P, Arora A, Strand TA, et al. Diagnostic accuracy of point of care tests for diagnosing celiac disease: a systematic review and meta-analysis. *J Clin Gastroenterol.* 2019;53:535–42.
9. Husby S, Koletzko S, Korponay-Szabó IR, et al. European Society for Pediatric Gastroenterology Hepatology, and Nutrition guidelines for the diagnosis of coeliac disease. *J Pediatr Gastroenterol Nutr.* 2012;54:136–60, <http://dx.doi.org/10.1097/MPG.0b013e31821a23d0>.
10. Lee GJ, Boyle B, Ediger T, et al. Hypertransaminasemia in newly diagnosed pediatric patients with celiac disease. *J Pediatr Gastroenterol Nutr.* 2016;63:340–3, <http://dx.doi.org/10.1097/MPG.0000000000001153>.

M.X. Espriu-Ramírez^{a,*}, Y. Rivera-Suazo^b, P.F. Valencia-Mayoral^c

^a *Gastroenterología y Nutrición Pediátrica, Hospital General de Cancún Dr. Jesús Kumate Rodríguez, Cancún, Quintana Roo, Mexico*

^b *Gastroenterología y Nutrición Pediátrica, Hospital Star Médica Infantil Privado, Mexico City, Mexico*

^c *Departamento de Patología Clínica y Experimental, Hospital Infantil de México Federico Gómez, Mexico City, Mexico*

* Corresponding author at: Professional Medical Center, piso 8, consultorio 812, Avenida Sayil SM 6, Manzana 5, Lote 2, Malecón Tajamar, Cancún, Quintana Roo, código postal 77500. Tel.: 998 898 1395.

E-mail address: draespriu@gmail.com (M.X. Espriu-Ramírez).

2255-534X/ © 2020 Asociación Mexicana de Gastroenterología. Published by Masson Doyma México S.A. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Monomorphic epitheliotropic intestinal T-cell lymphoma in a patient with chronic diarrhea and steatorrhea[☆]



Linfoma T intestinal monomórfico epiteliotrópico en un paciente con diarrea crónica y esteatorrea

Enteropathies are diseases that affect small bowel (SB) function and clinically manifest as chronic diarrhea and/or

steatorrhea.¹ They are a true clinical challenge, given their extensive differential diagnosis.

Monomorphic epitheliotropic intestinal T-cell lymphoma (MEITL) is a rare aggressive disease that mainly affects the SB mucosa. Its symptoms can be nonspecific, but often present as chronic diarrhea or steatorrhea. Diagnosis tends to be delayed, despite having distinctive histologic characteristics.

A 42-year-old man with alcohol use disorder presented with lenteric diarrhea and steatorrhea (on average 5–6 daily episodes), diffuse abdominal pain, edema, and weight loss of 20 kg, over a 22-month period. Physical examination revealed a cachectic aspect, ascites, and edema of the extremities. Laboratory test results showed severe malnutrition due to hypoalbuminemia (1.8 mg/dl) and anemia, with hemoglobin of 8.2 g/dl. HIV ELISA test was negative. Initial endoscopic studies were normal, and biopsies were taken of the ileum and colon. The biopsy of the distal ileum revealed villous atrophy of the mucosa and infiltra-

[☆] Please cite this article as: Ramírez-Quintero JD, Carvajal JJ, Echeverri C, Mosquera-Klinger G. Linfoma T intestinal monomórfico epiteliotrópico en un paciente con diarrea crónica y esteatorrea. *Revista de Gastroenterología de México.* 2021;86:319–322.




Figure 1 A) Segmental atrophy in the distal duodenum, B) Changes due to severe, homogeneous atrophy in the jejunum, C and D) Thinning of the mucosa in the mid jejunum with the presence of fibrin-covered superficial ulcerations in patches (blue arrows).




Figure 2 A) Hematoxylin and eosin $\times 20$: Dense lymphoid infiltrate composed of medium-sized monomorphic lymphoid cells. B, C, and D) Neoplastic cells express the CD3, CD8, and CD56 markers, respectively.

tion of small lymphocytes in the lamina propria. Given those histologic findings, celiac disease was suspected, and the patient was started on a gluten-free diet, but his clinical condition remained the same. Deamidated gliadin peptide antibody and tissue transglutaminase antibody tests were negative. Anterograde enteroscopy identified inflammatory

changes and villous atrophy in the distal duodenum and in the jejunum (Fig. 1A–D). Biopsies revealed a dense and monotonous infiltrate of small lymphocytes, with exocytosis into the epithelium and crypt destruction. Through immunohistochemistry, those lymphocytes expressed the CD3, CD7, CD8, and CD56 markers. They were negative for CD4 and

Table 1 Differential diagnosis in intraepithelial lymphocytosis associated with lymphoma.

Findings	Monomorphic epitheliotropic intestinal T-cell lymphoma	Enteropathy-associated T-cell lymphoma	Extranodal NK/T-cell lymphoma, nasal type
Endoscopic aspect	Inflammatory changes, atrophy, superficial ulcers	Severe inflammatory changes, villous atrophy, tumor-like appearance	Superficial ulcers, nodular or tumor-like appearance
Morphologic aspect	Monomorphic infiltrate of small-to-medium-sized lymphocytes	Polymorphic infiltrate of large lymphocytes, vesicular nuclei, prominent nucleoli, and scant cytoplasm	Polymorphic infiltrate of small, medium, and large lymphocytes; angiocentric pattern and extensive necrosis
Immunohistochemistry	CD3+, CD8+, CD56+	CD3+, CD8-/+ , CD56-	CD3-, CD8-, CD56+
Gastrointestinal dominant sites	Duodenum, jejunum	Proximal jejunum, rare in distal ileum and in colon	Nasopharynx, rare in stomach and jejunum
Associations of interest	No relation to enteropathy. Asians and Latin Americans.	Celiac disease. Frequent in white people.	High incidence in Asians. Related to Epstein-Barr virus.

NK: natural killer.

perforin and the CD5 T-cell marker was lost. Ki-67 showed high proliferative activity. Those findings are characteristic of MEITL (Fig. 2A–D). No adenopathy or organomegaly was reported in the staging studies, but the bone marrow was infiltrated by atypical lymphocytes possessing the same intestinal immunophenotype.

The patient’s progression during hospitalization was poor. He presented with Wernicke’s encephalopathy, gastrointestinal bleeding, and finally death, due to sepsis secondary to *Staphylococcus aureus* bacteremia, without having begun chemotherapy.

The differential diagnosis of enteropathy with steatorrhea is very broad and includes immune system-mediated causes, such as celiac disease, Crohn’s disease, and autoimmune enteropathy; drug-related causes; infiltrative causes, such as amyloidosis, eosinophilic enteritis, and collagenous sprue; infectious causes, such as HIV enteropathy, tropical sprue, giardiasis, Whipple disease, and tuberculosis; and rare causes, such as diffuse large B-cell lymphoma, enteropathy-associated T-cell lymphoma (EATL), and MEITL.¹ The presence of intraepithelial lymphocytosis significantly reduces the differential diagnosis to celiac disease, drug-induced enteropathy, EATL, and MEITL. Those diseases should be distinguished by immunohistochemical markers. The most useful in the differential diagnosis of lymphoma-associated intraepithelial lymphocytosis are CD3, CD8, and CD56 (Table 1).

T-cell lymphomas account for just 5% of gastrointestinal tract lymphomas (EATL and MEITL). EATL corresponds to nearly 90% of cases, is associated with celiac disease, tends to present in Whites, and is histologically characterized by a polymorphous infiltrate with numerous eosinophils and plasmacytes that can hide neoplastic lymphoid cells. MEITL is an aggressive pathology with poor outcome and predominates in Asian and Latin American males. Its median survival rate is usually 7 months.² At present, only one case has been reported in Colombia.³ The disease is not linked to celiac enteropathy, albeit a few cases have been described with said association.² Interestingly, in Asia, where celiac disease

is very rare, almost all cases of intestinal T-cell lymphoma are regarded as MEITL.⁴ Endoscopic findings of MEITL are inflammatory changes, such as edema, erythema, granularity, and superficial inflammatory ulcers.⁵ It spreads diffusely in the mucosa, with or without tumor-like lesions, unlike EATL, which is frequently associated with large tumors of the SB that tend to become perforated.² MEITL used to be considered a type II EATL, but its distinctive immunophenotypic and morphologic findings have enabled the clear differentiation between the two pathologies. In addition, the genomic hybridization technique frequently shows an 8q24 gain in MEITL, resulting in the definition of a different genomic profile.^{6,7}

There is no standard treatment for the disease. Multiple chemotherapy regimens have been utilized that are based on cyclophosphamide, doxorubicin, vincristine, and prednisolone (CHOP), PEG-asparaginase, pralatrexate, or autologous bone marrow transplantation, with varying efficacy.

In conclusion, MEITL is a rare entity, described in Asian and Latin American individuals, with a poor outcome and high mortality rate. Its clinical manifestations are diverse and its histopathologic findings distinctive, such as intraepithelial lymphocytosis with small-to-medium sized lymphocytes that are positive for CD3, CD8, and CD56. Those findings make the differential diagnosis of intraepithelial lymphocytosis a necessity in the clinical setting of chronic diarrhea and/or steatorrhea.

Ethical disclosures

The present work meets the current bioethics research regulations and was authorized by the institutional ethics committee. A written statement of informed consent was not requested, given that the data are carefully protected. No data from the clinical history or imaging studies could identify the patient described in the clinical case.

The authors declare that the present article contains no data that could identify the patient.

Financial disclosure

No financial support was received in relation to this article.

Conflict of interest

The authors declare that there is no conflict of interest.

References

- Jansson-Knodell CL, Hujoel IA, Rubio-Tapia A, et al. Not all that flattens villi is Celiac disease: a review of enteropathies. *Mayo Clin Proc.* 2018;93:509–17, <http://dx.doi.org/10.1016/j.mayocp.2017.10.025>.
- Lenti MV, Biagi F, Lucioni M, et al. Two cases of monomorphic epitheliotropic intestinal T-cell lymphoma associated with coeliac disease. *Scand J Gastroenterol.* 2019;54:965–8, <http://dx.doi.org/10.1080/00365521.2019.1647455>.
- Martínez-Baquero D, Galvis-Moreno L, Acosta-Forero J, et al. Linfoma T intestinal monomórfico CD56+. Primer caso informado en Colombia y revisión de la literatura. *Rev Fac Med.* 2015;63:127–31, <http://dx.doi.org/10.15446/revfacmed.v63n1.4746>.
- Yi JH, Lee GW, Do YR, et al. Multicenter retrospective analysis of the clinicopathologic features of monomorphic epitheliotropic intestinal T-Cell lymphoma. *Ann Hematol.* 2019;98:2541–50, <http://dx.doi.org/10.1007/s00277-019-03791-y>.
- Hashimoto R, Matsuda T. Gastrointestinal: endoscopic findings of monomorphic epitheliotropic intestinal T-Cell lymphoma (MEITL). *J Gastroenterol Hepatol.* 2018;34:311, <http://dx.doi.org/10.1111/jgh.14474>.
- Swerdlow SH, Campo CE, Harris NL, et al. WHO classification of tumours of haematopoietic and lymphoid tissues. *WHO Classification of Tumours, Volume 2, 4th Edition; 2017.*
- McElroy MK, Read WL, Harmon GS, et al. A unique case of an indolent CD56-positive T-cell lymphoproliferative disorder of the gastrointestinal tract: a lesion potentially misdiagnosed as natural killer/T-cell lymphoma. *Ann Diagn Pathol.* 2011;15:370–5, <http://dx.doi.org/10.1016/j.anndiagpath.2010.05.007>.

J.D. Ramírez-Quintero^a, J.J. Carvajal^b, C. Echeverri^c, G. Mosquera-Klinger^{b,*}

^a *Sección de Medicina Interna, Hospital Pablo Tobon Uribe, Medellín, Colombia*

^b *Unidad de Gastroenterología y Endoscopia Digestiva, Hospital Pablo Tobón Uribe, Medellín, Colombia*

^c *Hospital Pablo Tobón Uribe, Medellín, Colombia*

* Corresponding author at: Calle 78B #69-240, Consultorio 154, Hospital Pablo Tobón Uribe, Medellín, Colombia. E-mail address: gamkgastro@gmail.com (G. Mosquera-Klinger).

2255-534X/ © 2020 Asociación Mexicana de Gastroenterología. Published by Masson Doyma México S.A. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Splenic rupture after colonoscopy: A little-known complication[☆]



Rotura esplénica tras colonoscopia, una complicación poco conocida

Colonoscopy is a very frequently used diagnostic technique that has a good safety profile, but it is not exempt from complications, the most well-known of which are bleeding and perforation. However, there are other less common complications, such as splenic rupture, that have elevated morbidity and mortality.

A 77-year-old man had a history of GOLD III chronic obstructive pulmonary disease, lung adenocarcinoma with no current evidence of recurrence, atrial flutter (anticoagulation with edoxaban), prostatectomy due to adenocarcinoma, cholecystectomy, and right inguinal hernioplasty. He sought medical attention for diarrhea, asthenia, and weight loss of several months of progression. Upon admission, oral anticoagulation was suspended and substituted with enoxaparin (60 mg/12 h). Diagnostic colonoscopy was performed, with the patient under deep sedation with propofol. There

was no technical difficulty or immediate complications, and the only finding was diverticulosis. No pressure was applied on the left upper quadrant of the abdomen. Twelve hours after the procedure, the patient presented with low blood pressure, tachycardia, and generalized abdominal pain that was poorly controlled with opioids. The pain was described as progressively worsening over several hours. Emergency laboratory tests showed anemia of 8.4 g/dl (previously 12.9 g/dl) and an abdominal x-ray ruled out perforation. Emergency abdominal CT scan was carried out (Fig. 1)




Figure 1 Hemoperitoneum and a large peri-splenic hematoma measuring 10 × 12 × 13 cm (*) with foci of contrast medium extravasation in the lower splenic pole (white arrow), corresponding to a grade V splenic injury.

[☆] Please cite this article as: Rodríguez Prida J, Izquierdo Romero M, Álvarez Oltra G, López Caleyá JF, Ramírez Baum CE. Rotura esplénica tras colonoscopia, una complicación poco conocida. *Revista de Gastroenterología de México.* 2021;86:322–323.