



REVISTA DE GASTROENTEROLOGÍA DE MÉXICO

www.elsevier.es/rgmx



EDITORIAL

Evaluation of nutritional status and its association with the grade of activity of ulcerative colitis in Mexican patients[☆]



Evaluación del estado nutricional y su asociación con el grado de actividad en pacientes mexicanos con colitis ulcerosa crónica idiopática (CUCI)

Ulcerative colitis (UC) is a chronic and incurable disease that affects the mucosa and submucosa of the colon and rectum. Its etiology is unknown, but it is thought to be multifactorial, given that genetic, environmental, and immunologic factors are involved.¹ It is a well-known fact that chronic inflammation leads to chronic malnutrition due to protein catabolism, reduced protein synthesis, and fecal protein loss, as well as to an increase in bone resorption that produces the development of osteopenia and osteoporosis.^{2,3}

The presence of hypoalbuminemia has been evaluated as a predictive factor of severe activity and a risk factor for colectomy in Mexican patients with UC.⁴ Therefore, it is important to evaluate UC activity to induce or maintain remission and consequently improve the nutritional status of patients with UC. There are several scales for evaluating UC activity. The Truelove-Witts scale (TWS),⁵ Mayo endoscopic scale,⁶ and the Riley histologic index⁷ are among the most well-known and widely used. They evaluate clinical, endoscopic, and histologic aspects, respectively, but have not been validated. Even though there are other scales, none of them comprehensively assesses disease activity. A novel integral disease index for evaluating UC activity has now been validated in the Mexican population and encompasses clinical, biochemical, endoscopic, and histopathologic aspects of the disease, enabling a complete and comprehensive evaluation.⁸

In the current issue of the *Revista de Gastroenterología de México*, De León-Rendón et al.⁹ assessed the clinical utility of the Controlling Nutritional Status Score (CONUT) as

a method of nutritional screening and severity prediction in Mexican patients with UC in their cross-sectional study on 60 patients seen at a tertiary referral hospital in Mexico City. The most relevant findings of the study were that the CONUT score detected the risk for malnutrition in 93.3% of the patients, the risk was moderate in 43.3% of them, and no risk for malnutrition was detected in only 6.3% of the patients with UC. The cases with higher CONUT scores had a greater frequency of moderate-to-severe activity on the TWS, whereas those with lower CONUT scores had mild activity or no disease activity on the TWS, with statistical significance (6.84 ± 2.86 vs. 4.1 ± 3.02 ; $p = 0.009$). Likewise, higher CONUT scores were significantly associated with C-reactive protein (CRP) levels ≥ 45 mg/dL, compared with patients in whom that marker was not elevated (7.37 ± 2.45 vs. 4.97 ± 3.19 ; $p = 0.002$), as well as with an erythrocyte sedimentation rate (ESR) ≥ 30 mm/h (6.78 ± 2.78 vs. 4.60 ± 3.26 ; $p = 0.009$). Importantly, the CONUT score was not significantly associated with more relevant outcomes, such as mucosal healing assessed by the Mayo endoscopic subscore ($p = 0.35$) or histologic remission evaluated with the Riley histologic index ($p = 0.24$), most likely due to the small sample size. The authors also determined the diagnostic utility of the CONUT score and found that a score ≥ 6 points was associated with higher TWS scores, with an area under the curve (AUC) of 0.75, $p = 0.003$, sensitivity (S) of 64%, specificity (Sp) of 88%, positive predictive value (PPV) of 83% and negative predictive value (NPV) of 50%. Regarding biochemical markers, the CONUT score was associated with an ESR > 30 mm/h (AUC = 0.68, $p = 0.01$, S of 63%, Sp of 75%, PPV of 83%, and NPV of 50%) and CRP > 45 mg/dL (AUC = 0.65, $p = 0.04$, S of 67%, Sp of 64%, PPV of 60%, and NPV of 70%).

The CONUT score is based on the quantification of albumin, total cholesterol, and the absolute lymphocyte count and has shown its clinical utility for determining

☆ Please cite this article as: Yamamoto-Furusho JK. Evaluación del estado nutricional y su asociación con el grado de actividad en pacientes mexicanos con colitis ulcerosa crónica idiopática (CUCI). Revista de Gastroenterología de México. 2021;86:107–109.

malnutrition.¹⁰ It has also been used as a severity and mortality predictor in heart disease and gastrointestinal tumors.¹¹⁻¹⁵

In the article discussed herein, the clinical utility of the CONUT score was evaluated for the first time in Mexican patients with UC. The scale was shown to be a useful tool in detecting malnutrition but was only associated with clinical parameters determined by the TWS and biochemical parameters established through CRP levels and the ESR. Unfortunately, the CONUT score was not associated with endoscopic remission evaluated by the Mayo endoscopic subscore or histologic remission by the Riley histologic index, which are current aims in the medical treatment of patients with UC. Thus, the CONUT score is adequate for the nutritional evaluation of patients with UC, but it is not a useful tool for predicting severity, from an endoscopic or histologic perspective. Another aspect to take into account is that the evaluation of patients with UC should comprehensively include clinical, biochemical, endoscopic, and histopathologic aspects, as is the case with the novel integral disease index for evaluating the grade of activity of ulcerative colitis, or Yamamoto-Furusho index, that is now validated in the Mexican population,⁸ and whose aim is to optimize medical treatment and prevent adverse outcomes, such as surgery, hospitalization, and poor quality of life.

Finally, the evaluation of nutritional status through scales, such as the CONUT score, is important because it enables early nutritional intervention, which can prevent patients from progressing to a status of significant malnutrition. That is especially true for patients with clinically and biochemically determined moderate-to-severe activity, as demonstrated by the CONUT score.

Ethical considerations

No patients participated in the present study nor were patient data used, therefore no statements of informed consent were required. Likewise, given that there was no intervention, maneuver, or data management, the study is considered low-risk and no review or approval by the local ethics committee was needed. Even so, the study meets the current regulations regarding research, personal and identification data confidentiality, and participant anonymity (all healthcare workers that participated voluntarily). The present article contains no personal information that could identify the participants.

Funding

None.

Conflict of interest

Dr. Jesús Kazuo Yamamoto Furusho is a member of the advisory board, an opinion leader, and speaker for Abbvie Laboratories of Mexico, Abbvie International, Takeda International, Takeda Mexico, Pfizer (international and regional), and Janssen Cilag (international, regional, and national). He is an opinion leader and speaker for Alfasigma, Celltrion, Ferring, and Farmasa Schwabe, and has been a

speaker for Almirall, Astra Zeneca, Danone, and Novartis. He has been a research advisor for UCB Mexico and has received funding for research studies from the Shire, Bristol Myers Squibb, Pfizer, Takeda, and Celgene laboratories.

References

- Yamamoto-Furusho JK. Genetic factors associated with the development of inflammatory bowel disease. *World J Gastroenterol.* 2007;13:5594-7.
- Bueno-Hernández N, Núñez-Aldana M, Ascaño-Gutierrez I, Yamamoto-Furusho JK. Evaluation of diet pattern related to the symptoms of mexican patients with Ulcerative Colitis (UC): through the validity of a questionnaire. *Nutr J.* 2015; 14:25.
- Rodríguez-Bores L, Barahona-Garrido J, Yamamoto-Furusho JK. Basic and clinical aspects of osteoporosis in inflammatory bowel disease. *World J Gastroenterol.* 2007;13:6156-65.
- Choquet A, Yamamoto-Furusho JK, Reyes E, et al. Predictors of colectomy in patients with ulcerative colitis. A cohort analysis of 184 cases. *Rev Invest Clin.* 2004;56:11-5.
- Truelove SC, Witts LJ. Cortisone in ulcerative colitis; final report on a therapeutic trial. *Br Med J.* 1955;2:1041-8.
- Schroeder KW, Tremaine WJ, Ilstrup DM. Coated oral 5-aminosalicylic acid therapy for mildly to moderately active-ulcerative colitis. *N Engl J Med.* 1987;317:1625-9.
- Riley SA, Mani V, Goodman MJ. Microscopic activity in ulcerative colitis: what does it mean? *Gut.* 1991;32:174-8.
- Yamamoto-Furusho JK, Bozada-Gutiérrez KE, Sánchez-Rodríguez A, Bojalil-Romano F, Barreto-Zúñiga R, Martínez Benítez B. Validation of a novel integral disease index for evaluating the grade of activity in Mexican patients with ulcerative colitis: a prospective cohort study. *Rev Gastroenterol Mex.* 2019;84:317-25.
- De-León-Rendón JL, López-Pérez RY, Gracida-Mancilla NI, Jiménez-Bobadilla B, Alarcón-Bernes L, Mendoza-Ramírez S, et al. The controlling nutritional status score: a promising tool for nutritional screening and severity predictor in ulcerative colitis patients. *Rev Gastro Mex.* 2021:86.
- Ignacio de Ulíbarri J, González-Madroño A, de Villar NGP, González P, González B, Mancha A, et al. CONUT: a tool for controlling nutritional status, first validation in a hospital population. *Nutr Hosp.* 2005;20:38-45.
- Takehara K, Sakamoto K, Takahashi R, et al. Superior mesenteric artery syndrome improved by enteral nutritional therapy according to the Controlling Nutritional Status score. *Case Rep Gastroenterol.* 2017;11:729-35, <http://dx.doi.org/10.1159/000484129>.
- Sun X, Luo L, Zhao X, Ye P. Controlling Nutritional Status (CONUT) score as a predictor of all-cause mortality in elderly hypertensive patients: a prospective follow-up study. *BMJ Open.* 2017;7:e015649.
- Miyata T, Yamashita YI, Higashi T, Taki K, Izumi D, Kosumi K, et al. The Prognostic Impact of Controlling Nutritional Status (CONUT) in intrahepatic cholangiocarcinoma following curative hepatectomy: a retrospective single institution study. *World J Surg.* 2018;42:1085-91.
- Shirakabe A, Hata N, Kobayashi N, Okazaki H, Matsushita M, Shibata Y, et al. The prognostic impact of malnutrition in patients with severely decompensated acute heart failure, as assessed using the Prognostic Nutritional Index (PNI) and Controlling Nutritional Status (CONUT) score. *Heart Vessels.* 2018;33:134-44.
- Niwano M. The survival prognosis of elderly undernourished inpatients admitted to the internal medical department of an

emergency hospital as assessed using the nutritional screening, tool CONUT (for CONtrolling NUTritional status). *Nihon Ronen Igakkai Zasshi.* 2017;54:356–63.

J.K. Yamamoto-Furusho*

*Clinica de Enfermedad Inflamatoria Intestinal,
Departamento de Gastroenterología, Instituto Nacional de
Ciencias Médicas y Nutrición Salvador Zubirán, Mexico
City, Mexico*

*Corresponding author. Fundador y Director de la Clínica de Enfermedad Inflamatoria Intestinal, Departamento de Gastroenterología, Instituto Nacional de Ciencias Médicas y Nutrición Salvador Zubirán, Vasco de Quiroga 15, Colonia Belisario Domínguez, Sección XVI, Delegación Tlalpan, CP 14080 Ciudad de México, Mexico. Tel.: 5555733418; Fax: +5556550942.

E-mail address: kazuofurusho@hotmail.com