



ORIGINAL ARTICLE

Association between follicular gastritis and *Helicobacter pylori* in children seen at a public hospital in Peru[☆]



CrossMark

C.R. Mejia^{a,b,*}, C.A. Vera^{c,d}, L. Huiza-Espinoza^e

^a Escuela de Medicina Universidad Peruana de Ciencias Aplicadas-UPC, Lima, Peru

^b Asociación Médica de Investigación y Servicios en Salud, Lima, Peru

^c Universidad Ricardo Palma, Lima, Peru

^d Sociedad Científica de Estudiantes de Medicina de la Universidad Ricardo Palma, Lima, Peru

^e Servicio de Patología, Hospital Nacional Madre Niño «San Bartolomé», Lima, Peru

Received 2 August 2015; accepted 27 January 2016

Available online 27 April 2016

KEYWORDS

Helicobacter pylori;
Children;
Gastritis;
Peru (source: DeCS
BIREME)

Abstract

Background: For the last 15 years, infection from *Helicobacter pylori* (*H. pylori*) has been recognized in gastritis pathogenesis, and is known to trigger an important inflammatory response in these patients.

Aim: To determine the association between follicular gastritis and *H. pylori* infection in children seen at a public hospital in Peru.

Methodology: An analytic, cross-sectional study was conducted on all the children treated at the *Hospital Nacional Docente Madre "Niño San Bartolomé"* in Lima, Peru, within the time frame of 2011-2012. All the personal data from the patients' medical histories and endoscopic procedures were collected. The crude prevalence ratios (PR) were obtained and adjusted (aPR) with their 95% confidence intervals (95%CI), using generalized linear models with the binomial family and log link function.

Results: A total of 123 children met the study criteria. Forty-eight (39%) of the study sample were girls and the mean age of the children was 12 years. *H. pylori* was present in 44% of the sample and 9% presented with more than 100 bacteria per field (classified as +++) . Thirty-five percent of the children had esophagitis due to concomitant reflux. The presence of *H. pylori* was associated with follicular gastritis ($P < 0.01$; aPR: 2.3; 95% CI:1.49-3.49), adjusted by the children's age.

[☆] Please cite this article as: Mejia CR, Vera CA, Huiza-Espinoza L. Asociación entre gastritis folicular y *Helicobacter pylori* en niños atendidos en un hospital público peruano. Revista de Gastroenterología de México. 2016;82:80-85.

* Corresponding author. Av. Las Palmeras 5713-Los Olivos, Lima 39, Perú. Phone: +511 997643516.

E-mail addresses: christian.mejia.md@gmail.com, chrimeal@yahoo.com (C.R. Mejia).

Conclusions: Based on the data analyzed, it was concluded that the children with follicular gastritis had a greater likelihood of having *H. pylori* than those that did not present with gastritis. These results can be extrapolated to other similar populations and should be evaluated in each setting so that this does not become a public health problem within the next few years.

© 2016 Asociación Mexicana de Gastroenterología. Published by Masson Doyma México S.A. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

PALABRAS CLAVE

Helicobacter pylori;
Niños;
Gastritis;
Perú (fuente: DeCS
BIREME)

Asociación entre gastritis folicular y *Helicobacter pylori* en niños atendidos en un hospital público peruano

Resumen

Antecedentes: Durante los últimos 15 años la infección por *Helicobacter pylori* (Hp) ha sido reconocida en la patogénesis de la gastritis, desencadenando una importante respuesta inflamatoria.

Objetivo: Determinar la asociación entre gastritis folicular y la infección de Hp en niños atendidos en un hospital público peruano.

Metodología: Estudio transversal analítico desarrollado entre los años 2011 y 2012, con la totalidad de los niños atendidos en Hospital Nacional Docente Madre «Niño San Bartolomé», de Lima, Perú. Se recolectaron los datos personales de su historia clínica, datos del procedimiento endoscópico y sus hallazgos. Se obtuvieron las razones de prevalencia crudas y ajustadas (RPa) con sus intervalos de confianza del 95% (IC del 95%), usando los modelos lineales generalizados, con la familia binomial y la función de enlace log.

Resultados: Participaron 123 niños que cumplían con los criterios del estudio, 48 (39%) fueron niñas y la mediana de edad fue 12 años. Presentó Hp el 44% de ellos y el 9% presentó más de 100 bacterias por campo (clasificados como 3 cruces). El 35% tuvo esofagitis por reflujo concomitante. La presencia de Hp estuvo asociada a la gastritis folicular ($p < 0.01$; RPa: 2.3; IC del 95%, 1.49-3.49), ajustando por la edad de los niños.

Conclusiones: Sobre la base de los datos analizados se concluye que los niños con gastritis folicular tienen más probabilidad de tener Hp con respecto a los que no presentan gastritis. Los resultados pueden ser extrapolados a otras poblaciones similares, debiendo ser evaluado cada escenario para que esto no se convierta en un problema de salud pública en unos años.

© 2016 Asociación Mexicana de Gastroenterología. Publicado por Masson Doyma México S.A. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Helicobacter pylori (*H. pylori*) is a spiral Gram-negative microorganism that has been reported worldwide. Approximately one out of every 2 persons of the world population is infected with *H. pylori*.¹⁻⁴ For the last 15 years, *H. pylori* infection has been recognized in the pathogenesis of gastritis and peptic ulcer disease by triggering an important inflammatory response.^{5,6} The infection is acquired mainly in childhood and is associated with different gastrointestinal symptomatology, including recurrent abdominal pain, chronic hemorrhagic gastritis, and follicular gastritis, and it is sometimes asymptomatic.⁷⁻⁹ Recent studies suggest that chronic follicular gastritis in childhood can increase the risk for developing gastric neoplasia in adult life.¹⁰ In our environment, different studies have been conducted on the histopathologic characteristics of gastritis from *H. pylori*,¹¹ but whether there is a relation between that infection and its main characteristics, or if that relation exists in the pediatric population, has not been precisely determined.

Therefore, the aim of our work was to determine whether there was an association between follicular gastritis and *H. pylori* in children seen at a public hospital in the metropolitan area of Lima, Peru.

Methods

Study design and population

Evaluating the anatomopathologic registers of the *Hospital Nacional Docente Madre Niño San Bartolomé* of all the children that underwent esophagogastroduodenoscopy requiring gastric biopsy or biopsy of some section of the gastrointestinal tract between the years 2011 and 2012, an analytic cross-sectional study was conducted on those patients that had the possibility of being affected by *H. pylori*. The patients seen at a different hospital, those that did not have a gastric biopsy, or those whose medical records were incomplete, were excluded from the study (23 patients).

Non-probabilistic convenience sampling was carried out, calculating a minimum sample size of 97 patients utilizing a 10% expected error and 95% confidence interval.

Instruments and variables

A worksheet was designed that included the personal data from the medical history (age, sex) and data from the endoscopic procedure and its findings: year of evaluation, film number, sample location (esophagus, stomach and/or duodenum), and the anatomopathologic characteristics.

Chronic nonhemorrhagic gastritis was defined when there was gastritis with the presence of lymphoplasmacytic infiltrate (also known as round cell infiltrate) in the gastric mucosa. Chronic hemorrhagic gastritis was defined with the appearance of multiple erythematous and/or predominantly superficial ulcers at the corpus and antrum, and even reaching the duodenum. Follicular gastritis was defined as such when there were lymphocytes forming complete nodules and follicles. Reflux esophagitis was defined according to the consensus of the American Gastroenterology Association, using the pathologic criterion of abundant eosinophils (≥ 15) predominantly in the superficial layer of the epithelium, as well as the possibility of basal cell hyperplasia and elongated papillae.^{12,13} The dependent variable was the presence of *H. pylori*. In order to be observed and to evaluate its damage to the mucosa, it was classified as absent if no bacteria were found in the observation field, it was described with a plus sign (+) if 1-20 bacteria per field were observed, two plus signs (++) if 21-100 bacteria per field were observed, and three plus signs (+++) if more than 100 bacteria per field were observed. This classification has been used by national and international authors for diagnosing *H. pylori* in different types of samples.^{14,15} This process of *H. pylori* identification was aided by the use of hematoxylin and eosin staining and then Giemsa and Warthin Starry stains for confirming the presence of *H. pylori*. The histochemical staining was processed at the hospital in which the study was conducted, following the technical recommendations, and they were used as markers for evaluating the presence of *H. pylori*.

Procedures

Permission was requested from the baseline coordinators to collect the data coinciding with the study period. After permission was granted, coordination meetings were held to define the variables and aims. The missing information of the patients included in the final baseline was searched for in the hospital's database. And finally, the baseline was created, carrying out data quality control and analyzing the spreadsheet. The baseline was generated using the Windows Excel 2010 program and the data were processed using the Stata version 11.1 statistics software (StataCorp LP, College Station, TX, USA).

Data analysis

In the descriptive analyses, the numerical variables were evaluated with the normality suppositions and then

described through medians and interquartile ranges. The categorical variables were analyzed through percentages and frequencies. For the bivariate statistics, the Fisher's exact test was used for the cross-tabulation of the categorical variables and the rank sum test for the cross-tabulation of the categorical variables vs the quantitative ones, after the evaluation of the normality suppositions in each case. For the inferential statistics, generalized linear models were used, employing the binomial family and the log link function, obtaining the crude prevalence ratios and 95% confidence intervals (95% CI). The multivariate analysis was conducted in the same manner, obtaining the adjusted prevalence ratios (aPR). Statistical significance was set at a $p < 0.05$.

Ethical aspects

Data confidentiality was maintained and the ethical recommendations of the international organizations were followed. The study protocol was approved by the ethics committee of the *Hospital Nacional Docente Madre Niño San Bartolomé*.

Results

One hundred twenty-three patients that fit the study criteria were included. Of that total, 48 (39.0%) were girls. The median age was 12 years and the interquartile range was 7-14 years. Likewise, 73 (59.4%) biopsies were performed in the stomach and esophagus, 24 (19.5%) only in the stomach, 15 (12.2%) in the stomach, esophagus, and duodenum, 6 (4.9%) only in the esophagus, and 5 (4.1%) in the stomach and duodenum, according to the criteria of the specialist. The histopathologic characteristics in relation to the finding site are shown in Table 1.

Of the patient total 43.9% ($n=54$) were diagnosed with *H. pylori*, and of those patients, infection in 50% ($n=27$) of them was catalogued as + (1-20 bacteria per field), in 41% ($n=22$) as ++ (21-100 bacteria per field), and in 9% ($n=5$) as +++ (> 100 bacteria per field).

The number of plus signs varied according to whether the patient had follicular gastritis or not. A total of 55.6% ($n=15$) of those patients that had one plus sign, did not present with follicular gastritis. In contrast, 81.8% ($n=18$) of those that had 2 plus signs presented with follicular gastritis. And finally, 40.0% ($n=2$) of the patients that had 3 plus signs presented with follicular gastritis, and the differences were statistically significant ($p=0.02$).

The mean age varied depending on the type of pathology. The median age of those that presented with follicular gastritis was 13 years (interquartile range: 9-16 years) vs a median of 11 years (interquartile range: 4-14 years) of those with no follicular gastritis diagnosis ($p=0.01$). Likewise, the median age of those patients with *H. pylori* infection was 13 years (interquartile range: 10-15 years) vs a median of 9 years (interquartile range: 4-14 years) of those with no *H. pylori* diagnosis ($p < 0.01$).

The bivariate analysis showed a statistically significant relation between age ($p < 0.01$) and follicular gastritis ($p < 0.01$) with the presence of *H. pylori* in the children evaluated. The remaining results are shown in Table 2.

Table 1 Histopathologic characteristics according to the finding site in children infected with *Helicobacter pylori* at a public hospital in Peru.

Findings	n	%
<i>Stomach</i>		
Gastritis with follicles	44	37.6
Chronic nonhemorrhagic gastritis	31	26.5
Chronic hemorrhagic gastritis	20	17.1
Chronic gastritis in resolution with fibrin	2	1.7
Unaltered mucosa	8	6.8
Congestive mucosa	9	7.7
Hyperplasia or metaplasia	2	1.7
Adenocarcinoma	1	0.9
<i>Esophagus</i>		
Reflux esophagitis	26	27.4
No changes	14	14.6
Inflammatory or congestive	27	28.4
Acute esophagitis	9	9.5
Increased vasculature	8	8.4
Chronic esophagitis	9	9.5
Re-exacerbated chronic esophagitis	1	1.1
Gastric metaplasia	1	1.1
<i>Duodenum</i>		
Chronic duodenitis	16	80
Edema and infiltrate	3	15
Acute duodenitis	1	5.0

Table 2 Bivariate analysis of the presence of *Helicobacter pylori* according to characteristics of children seen at a public hospital in Peru.

Variable	p value	cPR	95%CI
Year of analysis, 2012	0.20	0.8	0.52-1.15
Male sex	0.47	0.9	0.58-1.29
Age (years)	<0.01	1.1	1.03-1.13
Nonhemorrhagic gastritis	0.30	0.8	0.45-1.28
Hemorrhagic gastritis	0.41	0.8	0.41-1.45
Gastritis with follicles	<0.01	2.6	1.75-3.89
Reflux esophagitis	0.08	1.4	0.95-2.16

cPR: crude prevalence ratio obtained through generalized linear models (GLM), using the binomial family and the log link function.

The multivariate analysis produced a statistically significant relation between follicular gastritis ($p < 0.01$) and the presence of *H. pylori* - adjusted by the children's age. **Table 3** shows the rest of the results.

Discussion

H. pylori infection has been studied from many perspectives, even though there is still much to discuss about this topic. However, there are clear concepts as to the anatomo-pathologic damage it can cause, but this has been studied very little in relation to childhood. It is known that a longer exposure time increases the risk for diseases caused by this pathology, and one of the factors to keep in mind - and the

Table 3 Multivariate analysis of the presence of *Helicobacter pylori* according to characteristics of the children seen at a public hospital in Peru.

	p value	aPR	95%CI
Age (years)	0.06	1.1	1.03-1.13
Follicular gastritis	<0.01	2.3	1.49-3.49

aPR: adjusted prevalence ratio obtained through generalized linear models (GLM) using the binomial family and the log link function.

most important in our work - is its relation to gastritis.^{16,17} Patients with follicular gastritis were found to have a 2.3-fold greater frequency of *H. pylori* infection, compared with those that did not have follicular gastritis. These results coincide with those of a similar study carried out in 2012 reporting that chronic gastritis, active gastritis, and follicular gastritis are significantly higher in children infected with *H. pylori*.¹⁸ Another similar study conducted in Colombia showed that the patients with this type of gastritis had more intense and persistent dyspeptic symptoms. In addition, it stated that there was substantial improvement of those symptoms and the histologic findings in the gastric mucosa upon *H. pylori* eradication.¹⁹ This is due to the fact that *H. pylori* induces the production of proinflammatory substances by mononuclear cells. In this manner, substances such as interleukin 12 create a greater stimulus for a greater presence of Th1 lymphocytes, forming lymphoid follicles, but they are incapable of eliminating the bacterium.

Even though age was not statistically significant in our study, this could have been due to the small sample size, given that the p value was very close to 0.05. Some studies have shown that there is an association between older age and *H. pylori* infection. A study carried out in Portugal stated that older children can contract *H. pylori* infection at a rate similar to that of small children.^{20,21} Another study conducted in Colombia demonstrated that *H. pylori* infection progressively increased with patient age and was more notable after 6 years of age.²² In Peruvian studies, the mean age for acquiring the infection has changed from 12-18 months in 1990 to 2.4 years in 2002.²³ This variable should be studied in future analyses, confirming whether it is an influential variable of the disease in our environment.

More than one fourth of the patients that presented with *H. pylori* also presented with reflux esophagitis. In a case-control study carried out in Brazil, it was found that reflux esophagitis was more frequent in children that did not have *H. pylori*, a result that differs from ours.²⁴ This could be due to the fact that their study population was one made up exclusively of esophageal biopsies, whereas our study also included gastric and duodenal biopsies. Further investigation is needed to determine whether there are differences in relation to the anatomic area evaluated.

The large majority of patients presented with chronic duodenitis, a result concurring with those of studies conducted in Brazil and Uganda. They reported that *H. pylori* infection was a risk factor for presenting with duodenal infections, especially if it was associated with *Giardia lamblia*.²⁵⁻²⁷ One out of every 10 patients presented with *H. pylori* infection described with 3 plus signs, meaning that

those children had more than 100 bacteria per field. Similar studies in Vietnam concurred that large numbers of *H. pylori* infection are associated with childhood.²⁸ Nevertheless, the quantity found in our study is alarming. This is due to the fact that we are a so-called Third World country, in which the prevalence of *H. pylori* infection is greater in the low economic population. *H. pylori* is also strongly associated with inadequate sanitary conditions in the developing countries.

And finally, a recent study stated that the diagnostic tests for defining treatment should be invasive (histology, antibiogram), leaving the noninvasive tests, as well as test/treatment, for epidemiologic or treatment confirmation studies, thus preventing unnecessary treatments and reducing strain resistance. Those children presenting with association with peptic ulcer disease should receive treatment adapted in accordance with sensitivity tests and lasting a recommended 10-14 days. Treatment costs, adherence, and possible adverse effects should be taken into account. A confirmation test, such as the breath test, is also recommended 4-8 weeks after treatment has ended.²⁹

A limitation of our study was selection bias, produced by not having an adequate sample, but even so, the data can be considered as an approximate reality value, given that they come from a referral hospital that treats a large number of children from the Peruvian capital. Moreover, there are no other studies in our environment that analytically assess the association between follicular gastritis and *H. pylori*, making ours an important starting point for future research on this topic. We recommend that further studies be conducted at more healthcare centers to evaluate aspects that were not analyzed in our study, as well as to determine whether age really is an influential factor in *H. pylori* infection.

Based on the data analyzed we conclude that pediatric patients with *H. pylori* infection have a higher frequency of follicular gastritis compared with those with non-*H. pylori* gastritis.

Ethical responsibilities

Protection of persons and animals. The authors declare that no experiments were performed on humans or animals for this study.

Data confidentiality. The authors declare that they have followed the protocols of their work center in relation to the publication of patient data.

Right to privacy and informed consent. The authors declare that no patient data appear in this article.

Financial disclosure

Self-financed.

Conflict of interest

The authors declare that there is no conflict of interest.

Acknowledgements

This article was prepared as part of the activities of the second Research Group of the SOCEM, offered in conjunction with the Medical Association of Research and Health Services. This manuscript was prepared by Claudia Vera, fulfilling the participation requirements of the second GIS, Lima, Peru.

References

- Garhart C, Czinn S. *Helicobacter pylori* infection: Review of pathogenesis and immunity. *Int Semin Paediatr Gastroenterol Nutr.* 2004;12:3-7.
- Drumm B, Day AS, Gold B, et al. *Helicobacter pylori* and peptic ulcer: Working Group of the Second World Congress of Pediatric Gastroenterology, Hepatology, and Nutrition. *J Pediatr Gastroenterol Nutr.* 2004;39:S626-31.
- Herbst JJ. Ulcer disease. In: Behrman RE, Kliegman RM, Jenson HB, editors. *Nelson textbook of pediatrics.* 16th ed. Philadelphia: Saunders; 2000. p. 1147-8.
- Go MF, Crowe SE. Virulence and pathogenicity of *Helicobacter pylori*. *Gastroenterol Clin North Am.* 2000;29:649-70.
- Go MF. Review article: Natural history and epidemiology of *Helicobacter pylori* infection. *Aliment Pharmacol Ther.* 2002;16:3-15.
- Makola D, Peura DA, Crowe SE. *Helicobacter pylori* infection and related gastrointestinal diseases. *J Clin Gastroenterol.* 2007;41:548-58.
- Raymond J, Bergeret M, Benhamou PH, et al. A 2-year study of *Helicobacter pylori* in children. *J Clin Microbiol.* 1994;32:461-3.
- Sbeih F, Abdullah A, Sullivan S, et al. Antral nodularity, gastric lymphoid hyperplasia, and *Helicobacter pylori* in adults. *J Clin Gastroenterol.* 1996;22:227-30.
- Das BK, Kakkar S, Dixit VK, et al. *Helicobacter pylori* infection and recurrent abdominal pain in children. *J Trop Pediatr.* 2003;49:250-2.
- Ladas SD, Rokkas T, Georgopoulos S, et al. Predictive factors and prevalence of follicular gastritis in adults with peptic ulcer and nonulcer dyspepsia. *Dig Dis Sci.* 1999;44:1156-60.
- Maghidman S, Cok J, Bussalleu A. Hallazgos histopatológicos en la gastritis nodular experiencia en el Hospital Nacional Cayetano Heredia. *Rev Gastroenterol.* 2001;21:261-70.
- Furuta GT, Liacouras CA, Collins MH, et al. Eosinophilic esophagitis in children and adults: A systematic review and consensus recommendations for diagnosis and treatment. *Gastroenterology.* 2007;133:1342-63.
- Villarín AJL, de Rezende L. Esofagitis eosinofílica. Revisión de los conceptos fisiopatológicos y clínicos actuales. *Gastroenterol Hepatol.* 2007;30:234-43.
- Bayona-Rojas MA, Gutierrez-Escobar AJ, Sánchez-Suarez JF, et al. Eficacia del método de inmunocromatografía en heces para el diagnóstico de *Helicobacter pylori* en pacientes con dispepsia: evaluación preliminar. *Respuestas.* 2014;19:79-85.
- Urribarri AM, García JC, Rivera AB, et al. *Helicobacter pylori* en niños atendidos en el Hospital Nacional Cayetano Heredia durante los años 2003 al 2006. *Rev Gastroenterol Perú.* 2008;28:109-18.
- Ramírez-Ramos A, Gilman R. *Helicobacter pylori* en el Perú. Lima-Perú: Editorial Santa Ana S.A; 2004.
- Marshall BJ, Warren JR. Unidentified curved bacilli in the stomach of patients with gastritis and peptic ulceration. *Lancet.* 1984;1:1311-4.
- Mazigh Mrad S, Abidi K, Brini I, et al. Nodular gastritis: An endoscopic indicator of *Helicobacter pylori* infection in children. *Tunis Med.* 2012;90:789-92.

19. Martínez Marín JD, Henao Riveros SC. Hiperplasia linfoides folicular gástrica e infección por *Helicobacter pylori* en adultos colombianos. Rev Col Gastroenterol. 2009;24:148-56.
20. Nguyen TV, Bengtsson C, Nguyen GK, et al. Age as risk factor for *Helicobacter pylori* recurrence in children in Vietnam. Helicobacter. 2012;17:452-7.
21. Oleastro M, Pelerito A, Nogueira P, et al. Prevalence and incidence of *Helicobacter pylori* infection in a healthy pediatric population in the Lisbon area. Helicobacter. 2011;16:363-72.
22. Gutiérrez O, Aponte D. Seroprevalencia y factores de riesgo asociados con la infección por *Helicobacter pylori* en niños. Rev Col Gastroenterol. 2001;16:19-22.
23. Mendoza D, Herrera P, Gilman RH, et al. Variation in the prevalence of gastric cancer in Perú. Int J Cancer. 2008;123:414-20.
24. Carvalho MA, Machado NC, Ortolan EV, et al. Upper gastrointestinal histopathological findings in children and adolescents with nonulcer dyspepsia with *Helicobacter pylori* infection. J Pediatr Gastroenterol Nutr. 2012;55:523-9.
25. Ankarklev J, Hestvik E, Lebbad M, et al. Common coinfections of *Giardia intestinalis* and *Helicobacter pylori* in non-symptomatic Ugandan children. PLoS Negl Trop Dis. 2012;6:e1780.
26. Escobar-Pardo ML, de Godoy AP, Machado RS, et al. Prevalence of *Helicobacter pylori* infection and intestinal parasitosis in children of the Xingu Indian Reservation. J Pediatr (Rio J). 2011;87:393-8.
27. Rodrigues MN, Queiroz DM, Bezerra Filho, et al. Prevalence of *Helicobacter pylori* infection in children from an urban community in north-east Brazil and risk factors for infection. Eur J Gastroenterol Hepatol. 2004;16:201-5.
28. Tindberg Y, Nyren O, Blennow M, et al. *Helicobacter pylori* infection and abdominal symptoms among Swedish school children. J Pediatr Gastroenterol Nutr. 2005;41:33-8.
29. Kalach N, Bontems P, Cadrelan S. Advances in the treatment of *Helicobacter pylori* infection in children. Annals of Gastroenterology: quarterly publication of the Hellenic Society of Gastroenterology. 2015;28(1):1-10.