



REVISTA DE GASTROENTEROLOGÍA DE MÉXICO

www.elsevier.es/rgmx



REVIEW ARTICLE

Dietary fiber and the microbiota: A narrative review by a group of experts from the *Asociación Mexicana de Gastroenterología*☆



A.T. Abreu y Abreu^a, M.P. Milke-García^b, G.A. Argüello-Arévalo^c,
A.M. Calderón-de la Barca^d, R.I. Carmona-Sánchez^e, A. Consuelo-Sánchez^f,
E. Coss-Adame^g, M.F. García-Cedillo^g, V. Hernández-Rosiles^f, M.E. Icaza-Chávez^h,
J.N. Martínez-Medinaⁱ, S. Morán-Ramos^j, E. Ochoa-Ortiz^k, M. Reyes-Apodaca^f,
R.L. Rivera-Flores^l, F. Zamarripa-Dorsey^m, F. Zárate-Mondragónⁿ, R. Vázquez-Frias^{f,*}

^a Hospital Ángeles Pedregal, Mexico City, Mexico

^b Dirección de Nutrición, Instituto Nacional de Ciencias Médicas y Nutrición Salvador Zubirán, Mexico City, Mexico

^c Departamento de Gastroenterología y Nutrición Pediátrica, Centro Médico Nacional La Raza, Instituto Mexicano del Seguro Social, Mexico City, Mexico

^d Departamento Nutrición y Metabolismo, Centro de Investigación en Alimentación y Desarrollo, Hermosillo, Sonora, Mexico

^e Práctica Privada, San Luis Potosí, S.L.P, Mexico

^f Departamento de Gastroenterología y Nutrición, Instituto Nacional de Salud Hospital Infantil de México Federico Gómez, Mexico City, Mexico

^g Departamento de Gastroenterología, Instituto Nacional de Ciencias Médicas y Nutrición Salvador Zubirán, Mexico City, Mexico

^h Hospital Faro del Mayab Star, Mérida, Yucatán, Mexico

ⁱ Unidad de Genómica de Poblaciones aplicada a la Salud, Instituto Nacional de Medicina Genómica, Mexico City, Mexico

^j Unidad de Genómica de Poblaciones aplicada a la Salud, Instituto Nacional de Medicina Genómica, Consejo Nacional de Ciencia y Tecnología, Mexico City, Mexico

^k Tecnológico de Monterrey, Mexico City, Mexico

^l Laboratorio de Investigación en Gastro-Hepatología, Instituto Mexicano del Seguro Social, Mexico City, Mexico

^m Departamento de Gastroenterología, Hospital Juárez de México, Mexico City, Mexico

ⁿ Departamento de Gastroenterología, Instituto Nacional de Pediatría, Mexico City, Mexico

Received 5 October 2020; accepted 12 February 2021

Available online 16 June 2021

KEYWORDS

Dietary fiber;
Gut microbiota;

Abstract Dietary fiber intake is one of the most influential and efficacious strategies for modulating the gut microbiota. Said fiber can be digested by the microbiota itself, producing numerous metabolites, which include the short-chain fatty acids (SCFAs). SCFAs have local and systemic functions that impact the composition and function of the gut microbiota, and

☆ Please cite this article as: Abreu y Abreu AT, Milke-García MP, Argüello-Arévalo GA, Calderón-de la Barca AM, Carmona-Sánchez RI, Consuelo-Sánchez A, et al., Fibra dietaria y microbiota, revisión narrativa de un grupo de expertos de la Asociación Mexicana de Gastroenterología, Rev Gastroenterol México. 2021;86:287–304.

* Corresponding author. Departamento de Gastroenterología y Nutrición, Hospital Infantil de México Federico Gómez, Doctor Márquez 162, Doctores, Cuauhtémoc, 06720, Mexico City, Mexico.

E-mail address: rovaf@yahoo.com (R. Vázquez-Frias).

Nutrition;
Gastrointestinal
diseases

consequently, human health. The aim of the present narrative review was to provide a document that serves as a frame of reference for a clear understanding of dietary fiber and its direct and indirect effects on health.

The direct benefits of dietary fiber intake can be dependent on or independent of the gut microbiota. The use of dietary fiber by the gut microbiota involves several factors, including the fiber's physiochemical characteristics. Dietary fiber type influences the gut microbiota because not all bacterial species have the same capacity to produce the enzymes needed for its degradation. A low-fiber diet can affect the balance of the SCFAs produced. Dietary fiber indirectly benefits cardiometabolic health, digestive health, certain functional gastrointestinal disorders, and different diseases.

© 2021 Asociación Mexicana de Gastroenterología. Published by Masson Doyma México S.A. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

PALABRAS CLAVE

Fibra dietaria;
Microbiota intestinal;
Nutrición;
Enfermedades
gastrointestinales

Fibra dietaria y microbiota, revisión narrativa de un grupo de expertos de la Asociación Mexicana de Gastroenterología

Resumen Una de las estrategias que más impacto y mayor eficacia tiene para la modulación de la microbiota intestinal es el consumo de fibra dietaria, que puede ser digerida por la propia microbiota generando numerosos metabolitos. Entre éstos, se encuentran los ácidos grasos de cadena corta (AGCC) con funciones tanto locales como sistémicas, que impactan en la composición y función de la microbiota intestinal y por lo tanto en la salud humana. El objetivo de esta revisión narrativa fue generar un documento que sirva como marco de referencia para conocer acerca de la fibra dietaria y sus efectos directos e indirectos.

Los beneficios directos de la ingestión de fibra dietaria, pueden ser dependientes o independientes de la microbiota intestinal. La utilización de la fibra dietaria por esta última, depende de varios factores y de sus características fisicoquímicas. La clase de fibra dietaria influye sobre la composición de la microbiota intestinal debido a que no todas las especies tienen la misma capacidad de producir enzimas necesarias para su degradación. El consumo de dietas con bajo contenido de fibra dietaria puede afectar el balance de los AGCC producidos. Los beneficios indirectos de la fibra dietaria impactan sobre la salud cardiometabólica, la salud digestiva, ciertos trastornos funcionales gastrointestinales y enfermedades diversas.

© 2021 Asociación Mexicana de Gastroenterología. Publicado por Masson Doyma México S.A. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

The gut microbiota is currently recognized as playing a relevant role in human health. Advances in its study have been made on how its composition can be modulated, as well as on the metabolic function of the different microbial species that colonize the gastrointestinal tract, to improve human health and potentially prevent or treat diseases in general.¹ Some strategies can modulate the microbiota, such as the use of probiotics, prebiotics, and even fecal transplantation. One of the simplest and most efficacious is the intake of dietary fiber that is metabolizable by the gut microbiota itself, producing metabolites that include short-chain fatty acids (SCFAs), such as acetate, propionate, and butyrate, with both local and systemic functions. Through its impact on the composition and function of the gut microbiota, dietary fiber influences human health in general.

In February 2020, the *Asociación Mexicana de Gastroenterología* convened a multidisciplinary group of 17 specialists (clinical nutritionists, chemists with master's

degrees in nutrition, gastroenterologists, and pediatric gastroenterologists and nutritionists) to previously review, and then have an in-person discussion about the scientific evidence on the role of dietary fiber in the digestive physiology and health, in general, of children and adults. The aim of the present narrative review was to produce a document that serves as a frame of reference for a clear understanding of dietary fiber and its direct and indirect effects on health.

Dietary fiber

Over time, there have been different definitions of dietary fiber, based on physiologic aspects or on methods for its analysis by the Association of Official Agricultural Chemists (AOAC). The Institute of Medicine, now known as the National Academy of Medicine and is part of the National Academies of the United States, proposed a definition of dietary fiber to distinguish endogenous fiber in foods, or

Table 1 Types of fiber, according to their functionality, and examples.

Type of fiber	Examples
Dietary fiber	Lignin, cellulose, pectin, gums, β -glucans, and modified starch
Soluble fiber	Pectin, gums, β -glucans, wheat dextrin, <i>Psyllium</i> , and inulin
Fermentable fiber	Pectin, guar gum, β -glucans, wheat dextrin, inulin
Viscous fiber	Pectin, β -glucans, some gums (e.g., guar), <i>Psyllium</i>
Functional fiber	Resistant dextrin, <i>Psyllium</i> , fructooligosaccharides, polydextrose, isolated gums, isolated resistant starch
Insoluble fiber	Cellulose, lignin, some pectin, and some hemicelluloses
Nonfermentable fiber	Cellulose, lignin
Nonviscous fiber	Polydextrose, inulin

Source: Slavin, 2013.⁷

“dietary fiber”, from the fiber that is extracted or synthesized, called “functional fiber”, the sum of which is total fiber.² In 2009, the Codex Alimentarius Commission defined dietary fiber as “carbohydrate polymers with 10 or more monomeric units, which are not hydrolyzed by the endogenous enzymes in the small intestine of humans”,³ and is the definition that we use in the present technical position paper. Depending on the regulations of each country, including oligomers with 3 to 9 monomeric units in the definition is suggested.⁴

Dietary fiber can be classified according to its nature or origin, colligative properties, and fermentability.⁵ With respect to its nature or origin, it is classified as dietary (intrinsic or intact, that is found in foods) or functional (extracted or synthetic). Regarding its colligative properties, the fiber’s chemical structure defines two characteristics related to its mechanisms of action. The first is solubility: fiber can be soluble (with different degrees of solubility) or insoluble in water. The second is its gel-forming capacity: it can form a viscous or nonviscous solution, with the viscous solutions classified as having low, medium, or high viscosity. Regarding fermentability, fiber can be nonfermentable, partially fermentable (semi-fermentable), or completely fermentable. The physicochemical properties of the different types of fiber are not exclusive. The main properties are the capacity to react with water, to have a colligative property (solubility and gel formation or increased consistency or viscosity), to ferment, and to chelate. All those physicochemical properties sustain the functions of fiber in the organism. Among the characteristics of soluble fiber is the fact that it is more viscous and fermentable than insoluble fiber, it undergoes few changes, and it has a mechanical effect. However, different types of fiber have different combinations of those properties, and consequently, their effects on humans are different. Likewise, the same food can contain varying quantities of different types of fiber.

Tables 1 and 2 provide examples of fibers that can be grouped according to their degree of solubility and fermentability into:

- a) Short-chain, highly fermentable, soluble fiber: it is made up of oligosaccharides, such as fructooligosaccharides (FOSs) and galactooligosaccharides (GOSs) that stimulate bifidobacteria production. It has a weak laxative effect and does not affect bowel transit time, although it produces much gas.
- b) Long-chain, highly fermentable, soluble fiber: it stimulates bacterial growth in general, has a weak laxative effect, does not affect bowel transit time, and produces a moderate quantity of gases.
- c) Mediumly fermentable, partially soluble fiber: it has a good laxative effect, accelerates bowel transit, stimulates bacterial growth in general, and produces a moderate quantity of gases.
- d) Slowly fermentable insoluble fiber: it has a good laxative effect, stimulates bacterial growth, and produces a moderate quantity of gases.
- e) Nonfermentable insoluble fiber: it has a good laxative effect, accelerates bowel transit, and only stimulates the growth of specific bacteria that degrade it, such as *Xylanibacter* and *Prevotella*.

Importantly, the fermentation of dietary fiber gradually supplies energy, but the magnitude of the supply depends on the kind of fiber and the type of microbiota of the individual.^{6–8}

In addition to the types of fiber described above, other substances function like dietary fiber. Those substances are synthetic carbohydrates, such as polydextrose, methylcellulose, carboxymethylcellulose, hydroxypropyl methylcellulose, curdlan, scleroglucan, and analogues. Some synthetic oligosaccharides, nonabsorbable polyols (sorbitol and mannitol), saponins, tannins, phytates, and substances of animal origin, such as chondroitin and chitosan, also act like fiber.^{9,10} Chitosan is a polysaccharide of natural origin composed of α -1,4-linked glucosamine residues and is a component of the exoskeleton of crustaceans and the cell walls of fungi.

Some types of fiber can have a prebiotic effect because, upon being fermented in the colon, they selectively promote the growth or activity of the microbiota, which has functional and beneficial effects for the host. However, not all dietary fiber is prebiotic.¹¹ The primary prebiotics utilized in clinical studies are FOSs, GOSs, transgalactooligosaccharides (TOSs), xylooligosaccharides (XOSs), isomaltooligosaccharides (IMOSs), lactulose, hemicellulose (from sprouted barley), and inulin. Inulin is a fructan, or fructosan, a polysaccharide mainly composed of fructose units.¹²

Dietary fiber intake

Dietary fiber intake recommendations are age-dependent. The daily requirement for children over one year of age is calculated as age plus 5 or 10 g, or as 0.5 g/kg/day, for children over 2 years of age.¹³ In adolescents and adults, the calculation is 14 g for every 1000 kcal.¹⁴ That quantity of

Table 2 Types of fiber, properties, and sources.

Structure	Description and properties	Sources
Cellulose	Cell wall Polysaccharide of up to 10,000 glucose units per molecule Insoluble and resistant to enzyme digestion in the gastrointestinal tract	25% of the fiber in grains and fruit 30% of the fiber in vegetables and nuts Much of the fiber in bran
Hemicelluloses	Cell wall Polysaccharides with monomers other than glucose, associated with cellulose Partially soluble or insoluble	30% of the fiber in vegetables, fruits, legumes, and nuts Cereals
Pectin	Cell wall and intracellular tissue of fruits and vegetables Polysaccharides with galacturonic acid and a variety of monosaccharides Soluble in hot water.	Fruits (quince, Mexican hawthorn, citrus fruits), 15-20% of the fiber in vegetables, legumes, and nuts; sugarcane and potato
Inulin β -glucans	Vegetable cell plasma Cell wall of grains Branched glucose polymers that enable their formation of viscous solutions	Oats and barley
Resistant starch	Starch and starch degradation products that are nonabsorbable, or physically or chemically modified	1: Legumes 2: Under-ripe bananas 3: Formed by cooling foods with precooked starch (potatoes)
Nondigestible oligosaccharides	Oligosaccharides with 3-9 monomeric units They can form by polysaccharide hydrolysis They can be fermented	Cereals and nuts Onions, garlic, artichokes, chicory
Synthetic compounds	Derived from cellulose (methylcellulose, hydroxypropyl methylcellulose) Polydextrose, 50% colonic fermentation and bulking and prebiotic properties Fermentation is difficult	Polydextrose: as an ingredient it confers bulk, replaces sugar, and provides texture
Gums and mucilage	Gums: hydrocolloids derived from plant exudates Mucilage: present in the cells of the outer layer of seeds (<i>Psyllium</i>)	Gum arabic, tragacanth, guar, and carob Alginates or seaweed extracts (agar, carrageenans) Mucilage: <i>Psyllium</i> .
Lignin	It is not a polysaccharide but are bound to hemicellulose	Celery or the outer layer of cereal grains
Others	Phytic acid (associated with fiber), tannins, phytosterols	Cereal grains

Adapted from Gray.⁸

dietary fiber is not scientifically sustained, and therefore is referred to as the "recommended daily intake". There is great variability among the different institutions that make the recommendation. In Mexico, there are no systematically obtained complete tables of the fiber content in foods. The present nutritional value tables that refer to fiber content take data from other tables (generally not Mexican ones), in which the type of chemical analysis for measuring fiber is not registered. It is presumed that dietary fiber intake is greater in rural diets than urban diets because of the proportion of cereals, legumes, fruit, and vegetables they contain. The 2012 National Survey on Health and Nutrition (ENSANUT, the Spanish acronym) reflected deficient fiber consumption beginning at early ages.¹⁵ The 2016 midway ENSANUT showed that children 24 to 59 months of age with

food insecurity ingested even less fiber than their counterparts that did not suffer from food insecurity. The daily fiber intake in adolescents was 23.7 g in males and 21.2 g in females, and in older adults was even less: 20.2 g in men and 17.9 g in women.¹⁶

The World Health Organization recommends several strategies for increasing dietary fiber intake. Standing out among them is the dietary orientation aimed at changing the behavior of persons with scant fiber intake or that have a health alteration that requires greater intake. Greater consumption of natural sources of fiber (fruits, vegetables, legumes, whole grains) is also suggested, as well as the use of fiber supplements, when the recommended quantity is not achieved through diet or if it could be beneficial for a specific health problem.¹⁷

The direct impact of fiber

Properties of dietary fiber that influence the gut microbiota

The use of dietary fiber by the gut microbiota depends on its source, type of molecules, bonds, chain length, particle size, and association with other compounds.^{18,19} It also depends on the abovementioned physicochemical properties of solubility, viscosity, and fermentability.¹ The size of the particle determines its susceptibility to digestion, binding, water retention, and bowel transit time.²⁰ Water retention can influence the capacity of bacteria to infiltrate and digest fiber, as well as the speed of transport through the intestine.²¹ Viscosity depends on the degree of hydration, particle size, and the pH.²² Highly fermentable fiber, of which the β -glucans and pectins stand out, can also have high solubility and viscosity. The majority of soluble fibers are very viscous in the intestine. FOSs and pectins can be metabolized by bacteria in the ileum and ascending colon, unlike insoluble fiber, such as cellulose and hemicellulose, that is exclusively metabolized in the distal colon.¹

Experimentally, soluble fiber, compared with insoluble fiber, has been shown to modify the colonization of intestinal bacteria, impacting the richness of the gut microbiota.²³ Insoluble fiber intake produces a greater relative abundance of Bacteroidetes, Euryarchaeota and *Ruminococcaceae*, and at the genus level, of *Prevotella*, *Phascolarctobacterium*, *Coprococcus*, and *Leeia*. In contrast, soluble fiber intake produces a greater relative abundance of the phylum Proteobacteria and less abundance of *Prevotellaceae*, and with respect to genera, a greater abundance of *Blautia*, *Solobacterium*, *Syntrophococcus*, *Weissella*, *Olsenella*, *Atopobium*, and *Succinivibrio*.²⁴ Thus, in a study with 7% pectin, a soluble fiber, there was an increase in *Anaeroplasma*, *Anaerostipes*, and *Roseburia*, whereas there was a decrease in *Alistipes* and *Bacteroides* spp.¹⁷

Some of the components of fiber, such as the arabinoxylan oligosaccharides, can increase the abundance of bifidobacteria in the ascending colon, lactobacilli in the ascending colon and transverse colon, and *Clostridium coccoides* and *Eubacterium rectale* in the descending colon.¹⁷ The viscosity of fiber increases the number of anaerobic bacteria and *Clostridium* spp., whereas the number of aerobic bacteria and the genus *Enterococcus* negatively correlate with viscosity.²² Inulin fermentation results in a greater proportion of lactobacilli and bifidobacteria, a lower proportion of *Enterobacteriaceae*, and greater butyrate production.²⁴

The consumption of 12 g of inulin for 4 weeks in healthy adults with mild constipation induced an increase in the abundance of *Bifidobacterium* and *Anaerostipes* spp. and a decrease in the population of *Bilophila*.²⁵ Said effect could be attributed to the capacity of the genus *Bifidobacterium* to efficiently degrade FOSs and to the fact that *Anaerostipes hadrus* produces butyrate.²⁶ A similar effect occurred in patients with active ulcerative colitis that received 7.5 or 15 g of inulin enriched with oligofructose for 9 weeks, with an increase in the abundance of *Bifidobacteriaceae* and *Lachnospiraceae*, and an increase in butyrate production.²⁷

Resistant starch, which is a nondigestible fraction of cornstarch, raw potatoes, or unripe bananas, is also considered a dietary fiber. Its consumption is associated with an increase in the butyrate producers, *Ruminococcus bromii*, *Faecalibacterium prausnitzii*, and *E. rectale*. *R. bromii* is a key species for the fermentation of starches in the colon, as has been demonstrated in other studies.^{28,29}

Fiber and short-chain fatty acid (SCFA) production

Anaerobic bacteria in the large intestine produce SCFAs, through dietary fiber fermentation.³⁰ The principal SCFAs are acetate, propionate, and butyrate, at a ratio of 60:20:20. Lactate is the salt of a very common organic acid in the intestinal lumen that is also produced by bacteria, whereas other types of bacteria metabolize acetate, propionate, and butyrate. Thus, there are species that can be primary degraders of dietary fiber, whose products are degraded by other fermenting microorganisms that finally produce acetate, propionate, and butyrate.^{31,32}

Acetate is mainly produced via acetyl-CoA. Propionate is synthesized via two pathways: the succinate pathway and the acrylate pathway, from hexose and pentose or lactate substrates and via the propanediol pathway that utilizes deoxyhexoses, such as fucose and rhamnose, as substrates.³¹ The Bacteroidetes phylum produces propionate, mainly through the succinate pathway. Butyrate can also be produced from peptides or amino acids, and not only derived from dietary fiber sources. Some butyrate-producing species correspond to the families of *Ruminococcaceae* and *Lachnospiraceae*, both Firmicutes, as well as to *Erysipelotrichaceae* and *Clostridiaceae*. In addition, *F. prausnitzii* (the family *Ruminococcaceae*) can utilize polysaccharides that come from starch, hemicellulose, inulin, and pectin, whereas *E. rectale* is able to utilize starch, arabinoxylans, and inulin to produce SCFAs.³⁰

SCFAs perform different functions: they regulate both gene expression, by acting as inhibitors of histone deacetylases, and energy metabolism. They also act as signaling molecules that recognize specific receptors, thus promoting the regulation of the immune system and inflammation.^{30,31} The function of SCFAs varies, depending on the receptors in the host tissue that they can be assimilated in, giving rise to different physiologic effects.³² They carry out part of their function upon binding to the G protein-coupled receptors (GPCRs), also known as free fatty acid receptors, GPCR41 (or FFAR3), GPCR43 (or FFAR2), and GPCR109A. Acetate and propionate are potent GPCR43 activators, which are mainly expressed in colonocytes, adipose tissue, immune system cells, nervous system cells, and pancreatic cells and they are co-expressed with GLP-1 in the enteroendocrine cells. Thus, they are related to lipid or glucose metabolism, as well as to the immune system response.^{30,31}

A diet that is low in dietary fiber can affect SCFA production. In contrast, dietary fiber intake, and its effect on SCFA synthesis, can stimulate the production and secretion of intestinal mucus. That substance, which protects the intestinal mucosa, arises due to the increase in bacteria that promote gene expression in calciform cells or to the mechanical stimulus of dietary fiber itself.^{33,34}

Dietary fiber and its relation to the brain-gut-microbiota axis (BGMA)

There are different communication pathways in the BGMA. The main pathway is the vagus nerve, followed by enteric nervous system activity, upon producing molecules that act as neurotransmitters, such as gamma amino butyric acid (GABA), serotonin, melatonin, histamine, and acetylcholine. One of the main mechanisms that relates dietary fiber to the BGMA results from the direct influence of SCFAs and lactic acid, which also participate in the modulation of serotonin secretion.³⁵ Butyrate is primarily produced from resistant starch. Because one of the functions of butyrate is to inhibit the histone deacetylases, it has been shown to be beneficial in different neurologic diseases, such as Parkinson's disease, and to improve learning and memory in cases of dementia, including Alzheimer's disease, depression, and addictions. It is also thought to be a substrate for the production of energy in the brain, although it is not known to what degree. Cerebral inflammation has been reported to decrease in in vitro and in vivo models of Parkinson's disease.³⁶

FOSs and GOSs increase brain-derived neurotrophic factor (BDNF) gene expression, the NR1 and NR2A subunits of the N-methyl-D-aspartate (NMDA) receptors, and plasma peptide YY.³⁷ A study on mice showed that the microbiota modulated behavior, after recolonization of the mice with low concentrations of BDNF resulted in behavioral alterations.³⁸ GOSs suppress the response to neuroendocrine stress through the hyposecretion of cortisol and increase the attentional vigilance to positive stimuli versus negative stimuli.³⁹

The pharmacologic administration of sodium butyrate has been shown to have antidepressive effects.⁴⁰ Some bacteria, such as *Lactobacillus* and *Bifidobacterium* spp., are associated with neurologic development, emotional responses, and GABA production. *Streptococcus*, *Escherichia*, and *Enterococcus* spp. are associated with serotonin synthesis and *Bacillus* spp. is involved in dopamine production.⁴¹ Likewise, changes in BDNF and NMDA concentrations due to bacterial metabolism can contribute to the structural and chemical imbalances associated with schizophrenia and other psychopathologies.^{42–44}

Direct effects of dietary fiber on the diversity and abundance of the microbiota

One of the characteristics of the gut microbiota most consistently associated with a better health status is bacterial diversity.⁴⁵ Said diversity is importantly affected when diets are low in fiber or carbohydrates that are available for the microbiota.⁴⁶ The kind of dietary fiber influences the type of microbiota because not all species produce the enzymes necessary for fiber degradation.¹⁸

Different members of the gut microbiota are important dietary fiber degraders and include 130 glycoside hydrolases, 22 polysaccharide lyases, and 16 families of esterases, providing the flexibility to degrade different energy sources from the available fibers. The main species responsible for dietary fiber degradation are the phyla Firmicutes and Actinobacteria.^{34,47}

In a Finnish study on pregnant women with overweight or obesity, the consumption of whole grains and vegetables correlated with diversity of the microbiota. The quality of diet, in general, in the same study, correlated with abundance of the genus *Coprococcus* of the family *Lachnospiraceae*, the species *F. prausnitzii* of the family *Ruminococcaceae*, and an unknown species of the family *Barnesiellaceae*.⁴⁸ In general, differences in gut microbiota diversity between Western populations (Europe and the United States) and populations from Africa and Papua New Guinea have been demonstrated. Their diets also differ, with Western diets consisting of a high content of processed foods, meats, sugars, and saturated fats, whereas the African diet, especially in rural zones, is characterized by important consumption of vegetables, fruits, and whole-grain cereals.^{6,49–52}

In addition to the bacterial abundance of the microbiota, the different metabolites it produces are determinants of the physiologic effects on the host.⁵³ The consumption of soluble fiber, such as pectin and inulin, and insoluble fiber, such as hemicellulose, can increase the abundance of SCFA-producing bacteria.^{47,54–56} Butyrate is the main source of energy of the colonocytes and commensal bacteria, such as *E. rectale* and *F. prausnitzii*, participate in its production. That fatty acid participates in the regulation of proinflammatory and anti-inflammatory mechanisms.⁵⁷ Under said premise, butyrate has been reported to induce malignant cell apoptosis, thus reducing the risk for colorectal cancer.⁵⁸ Table 3 lists the types of dietary fiber, the food source, and the changes produced in the gut microbiota and in SCFA production.^{27,55,59–79} Importantly, not all individuals produce the same quantities of SCFAs, upon consuming different sources fiber, because of the varying capacity of the intestinal bacteria to produce those metabolites.

Microbiota and gut health

Specialized bacteria, such as *Clostridium*, *Lactobacillus*, and *Enterococcus*, capable of adhering to the gastrointestinal mucosa, feed on mucus and bond to epithelial cells.⁸⁰ Those types of bacteria have an important influence on the immune system and intestinal homeostasis.⁸¹ The gut microbiota also plays a significant role in maintaining mucosal integrity.⁴⁷ In germ-free experimental models, mucin-producing calciform cells decrease and the mucosal layer subsequently becomes thin.^{47,82} *Akkermansia muciniphila* has been identified as the determinant in maintaining that mucosal barrier,⁸³ *Bacteroides thuringiensis* has been reported to produce a strong bacteriocin against *Clostridioides difficile*, and *Bacteroides thetaiotaomicron* has been shown to participate in metalloproteinase expression, for the conversion of prodefensin into defensin.^{84,85} The microbiota also participates in the maintenance of Paneth cell integrity, and consequently, in the correct production of antimicrobial peptides. The increase in taxa, such as *Enterobacteriaceae*, *Pasteurellaceae*, *Veillonellaceae*, and *Fusobacteriaceae*, and the decrease in Erysipelotrichales, Bacteroidales, and Clostridiales, produce defects in the formation of those antimicrobial peptides, which is associated with inflammatory bowel disease (IBD).⁸⁶ There is an increase in Proteobacteria and a decrease in Firmicutes in

Table 3 Types of dietary fiber, food source, and changes induced in the gut microbiota and in short-chain fatty acid production.

Type of fiber, dose	Classification and food source	Changes in bacteria and their metabolites	Reference
Arabinoxyloligosaccharides 2.2 g/d	Soluble fiber Cereals	Increase in lactobacilli and <i>Bacteroides</i> Increase in butyrate	Walton et al., 2012 ⁵⁹
Arabinoxyloligosaccharides 3 g/d y 10 g/d	Soluble fiber Cereals	Increase in <i>Bifidobacterium</i> * Decrease in pH Increase in SCFAs	Francois et al., 2012 ⁶⁰
Arabinoxylans	Soluble fiber Cereals	Decrease in <i>Clostridium</i> clusters I/XI/XV and <i>Verrucomicrobia</i> Increase in Actinobacteria	Van den Abbeele et al., 2011 ⁶¹
Arabinoxyloligosaccharides 5 g/d	Soluble fiber Cereals	Increase in <i>Bifidobacterium</i> Decrease in isobutyric acid and isovaleric acid	Francois et al., 2014 ⁶²
β -glucans	Soluble and viscous fiber Oats, barley, wheat, rye, seaweed, mushrooms	Increase in <i>Bacteroides/Prevotella</i> , lactobacilli, bifidobacteria	Hughes et al., 2008 ⁵⁵ Snart et al., 2006 ⁶³ Jayachandran et al., 2018 ⁶⁴
Galactooligosaccharides 5.5 g/d	Soluble and nonviscous fiber Beans, chickpeas, lentils, onion, lettuce, broccoli, artichoke	Increase in <i>Bifidobacterium</i> spp. and <i>Bacteroides</i> spp. Changes in immune markers	Vulevic et al., 2015 ⁶⁵
Inulin	Soluble and nonviscous fiber Agave, artichoke, asparagus, banana, chicory root, onion, leeks, wheat	Increase in <i>Bacteroides</i> , <i>Lactobacillus</i> , <i>Bifidobacterium</i> , <i>Clostridium</i> , and <i>Lachnospiraceae</i> and decrease in enterobacteria	Valcheva et al., 2019 ²⁷
Agave inulin 5.0 and 7.5 g/d	Soluble and nonviscous fiber Agave and byproducts	Increase in butyrate Increase in <i>Bifidobacterium</i> Decrease in <i>Ruminococcus</i> , <i>Lachnobacterium</i> , <i>Desulfovibrio</i>	Holscher et al., 2014 ⁶⁶
Inulin + oligofructose 16 g/d	Soluble and nonviscous fiber Chicory	Increase in <i>Bifidobacterium longum</i> , <i>Bifidobacterium</i> <i>pseudocatenulatum</i> , and <i>Bifidobacterium</i> <i>adolescentis</i> Decrease in acetate and propionate	Salazar et al., 2015 ⁶⁷
Inulin + partially hydrolyzed guar gum 15 g/d	Soluble and nonviscous fiber/Soluble and viscous fiber	Decrease in <i>Clostridium</i> spp. No significant changes in SCFAs	Linetzky et al., 2012 ⁶⁸
Xyloligosaccharides 1.4 and 2.8 g/d	Soluble fiber	Increase in <i>Bifidobacterium</i> , <i>Bacteroides fragilis</i> No significant changes in pH, SCFAs, or lactic acid	Finegold et al., 2014 ⁶⁹
Polydextrose 8 g/d	Soluble and nonviscous fiber	Decrease in <i>Clostridium histolyticum</i> , lactobacilli/enterococci Increase in bacterial diversity No significant changes in SCFAs	Costabile et al., 2012 ⁷⁰

Table 3 (Continued)

Type of fiber, dose	Classification and food source	Changes in bacteria and their metabolites	Reference
Polydextrose 21 g/d	Soluble and nonviscous fiber	Increase in the Bacteroidetes: Firmicutes ratio Increase in <i>Parabacteroides</i> Decrease in <i>Eubacterium</i> , <i>Ruminococcus</i> , <i>Roseburia</i> , <i>Dorea</i> Decrease in fecal butyrate, phenol, and indole	Holscher et al., 2015 ⁷¹
Pectin	Soluble and viscous fiber Apples, cherries, oranges, carrots, apricots	Controversial, with respect to <i>Bacteroides</i> spp.	Dongowski et al., 2002 ⁷² Licht et al., 2010 ⁷³
Cornstarch 10 and 20 g/d	Soluble and nonviscous fiber Corn	Increase in <i>Parabacteroides</i> , <i>Bifidobacterium</i> Decrease in <i>Anaerostipes</i> , <i>Dorea</i> , <i>Ruminococcus</i> Decrease in fecal pH fecal and increase in SCFA production	Whisner et al., 2016 ⁷⁴
Cornstarch 40 g/d	Soluble and nonviscous fiber Corn	Increase in <i>Clostridium coccooides</i> , <i>Clostridium leptum</i> , <i>Lactobacillus</i> spp., <i>Parabacteroides distasonis</i> , and <i>Ruminococcus bromii</i> Decrease in <i>Ruminococcus gnavus</i> , <i>Ruminococcus</i> <i>torques</i> , and <i>Escherichia coli</i> Increase in SCFA production	Le Leu et al., 2015 ⁷⁵
Resistant starch 22-29 g/d	Soluble and nonviscous fiber Green bananas, potatoes, oats, beans	Increase in <i>Oscillospira guilliermondii</i> , <i>R. bromii</i> , <i>Sporobacter termitis</i> , <i>C. leptum</i> , <i>Clostridium</i> <i>cellulosi</i> , <i>Alistipes</i> spp., <i>Eubacterium rectale</i> Decrease in <i>Papillibacter cinnamivorans</i> and the microbiota diversity	Salonen et al., 2014 ⁷⁶
Cornstarch (resistant starch type 2)	Soluble fiber Corn	Increase in SCFAs Increase in <i>R. bromii</i> and <i>E. rectale</i> , <i>Faecalibacterium prausnitzii</i> , <i>Roseburia faecis</i> , and <i>Akkermansia muciniphila</i>	Martinez et al., 2010 ⁷⁷
Potato starch (resistant starch type 2)	Soluble fiber Potatoes	Increase in <i>B. adolescentis</i> and <i>R. bromii</i>	Martinez et al., 2010 ⁷⁷
Sweet potato residue	Soluble and insoluble fiber Sweet potatoes	Increase in <i>Bifidobacterium</i> and <i>Lactobacillus</i> and decrease in <i>Enterobacillus</i> , <i>Clostridium perfringens</i> , and <i>Bacteroides</i>	Martinez et al., 2010 ⁷⁷
Fructans	Soluble fiber Garlic, onion, wheat, rye, agave	Increase in the <i>Bifidobacterium</i> /enterobacteria ratio Increase in bifidobacteria and lactobacilli, <i>Eubacterium</i> , <i>Roseburia</i> , and <i>Faecalibacterium</i>	Ampatzoglou et al., 2015 ⁷⁸
Wheat bran fiber	Insoluble and nonviscous fiber Wheat bran	Increase in bifidobacteria, lactobacilli, <i>Atopobium</i> , enterococci, <i>Bacteroides</i> , and <i>Prevotella</i> Increase in butyrate	Freeland et al., 2009 ⁷⁹

SCFAs: short-chain fatty acids.

the different entities in which there is inflammation of the mucosa and an increase in permeability.^{87,88}

Indirect impact of fiber

Dietary fiber in cardiometabolic health

Diabetes mellitus (DM), obesity, dyslipidemia, high blood pressure, and metabolic syndrome are the most frequent cardiometabolic risk factors. Their appearance implies risks for future complications and even death, mainly when they present in a combined form. A proper diet contributes to maintaining health and a diet that is not adequate, balanced, or diverse is associated with the appearance of cardiometabolic alterations from infancy to adulthood.^{89,90} Less healthy dietary patterns, characterized by including little dietary fiber, correlate with greater cardiovascular risk.^{91,92} Two meta-analyses based on cohort studies found a lower mortality rate due to cardiovascular causes, in individuals that consume more dietary fiber.^{93,94}

Dietary fiber intake can reduce postprandial glycemia,⁹⁵ improve serum lipid levels,⁹⁶ and prevent obesity and the accumulation of visceral fat.⁹⁷ Dietary fiber, upon being metabolized by the gut microbiota, has been described to produce substrates that have a positive impact on the health of the host, particularly regarding cardiovascular health, because they reduce the risk for cardiovascular diseases and diabetes.^{98–100} In a systematic review and meta-analysis, from 19 studies, a lower risk for cardiovascular disease and coronary disease was associated with greater intake of total fiber, insoluble fiber, and fiber from cereals and vegetables. For each 7-gram increase of fiber intake per day, there was a RR of 0.91 (95% CI: 0.87–0.94) for coronary disease, as well as a reduced mortality rate (RR 0.59, 95% CI = 0.44, 0.78).⁹⁸ In another similar study conducted by the American Society for Nutrition, fiber intake was associated with a low and intermediate risk for cardiovascular disease and diabetes.¹⁰¹ Likewise, fiber intake reduced total cholesterol, LDL cholesterol, and triglycerides,¹⁰² as was reported in vegans and vegetarians, who showed a better lipid profile than individuals that ate less fiber and more meat.¹⁰³

Type 2 DM is associated with a decrease in bacteria that degrade dietary fiber. In studies utilizing animal models, the administration of soluble fiber, such as oligofructose and long-chain inulin, corrected the altered microbiota, or dysbiosis, reduced weight gain and low-grade inflammation, and improved glucose metabolism, intestinal permeability, and endotoxemia, partly related to the pathophysiology of DM.¹⁰⁴ In a 16-week study that included a diet supplemented with functional fibers, there was improvement in the colonic microbiota, characterized by a significant increase in bifidobacteria, lactobacilli, and *Bacteroides* counts, as well as a decrease in the clostridia count, with a decrease in LDL cholesterol and total cholesterol.¹⁰⁵ A rice bran extract induced the same decrease in postmenopausal women, in addition to reduced levels of TNF- α .¹⁰⁶ In individuals with hyperlipidemia, a plant-based diet reduced blood pressure and LDL cholesterol.¹⁰⁷ A decrease in inflammation (measured by C-reactive protein), overall inflammation, and cardiovascular risk has also been reported.¹⁰⁸

The French Nutrition and Health Survey concluded that dietary fiber and whole grain intake was inversely associated with systolic blood pressure.¹⁰⁹ The consumption of foods with a high glycemic index confers a greater risk for developing DM, compared with the effect of habitual dietary fiber and cereal intake.¹¹⁰ In fact, fiber intake ≥ 20 g/day reduces the risk for presenting with DM, most likely due to its effect on the proinflammatory state.¹¹¹ Other mechanisms for reducing the risk of DM have been posited, such as the adsorption of glucose by fiber in the gastrointestinal tract, slower gastric emptying, and improved postprandial insulinemia. However, not all intervention studies have shown beneficial results,¹¹² and the primary preventive effect of fiber in cardiovascular health has not been so obvious. That is possibly due to the fact that there are multiple genetic and environmental factors that are difficult to control, as well as to the small number of intervention studies, with discrepancies in the definitions employed, short intervention duration, and difficulty in conducting them.¹¹³ The conclusions of the most recent reviews and expert opinions on the topic coincide in underlining the fact that more studies with better quality methodologies are required.¹¹⁴

Dietary fiber, microbiota, and obesity

As mentioned above, dietary fiber intake can help prevent weight gain, visceral fat accumulation, and obesity.⁹⁷ Fiber intake is associated with other beneficial lifestyle factors, such as the consumption of fruits and vegetables and exercise habits. High-fiber diets are typically lower in fat and energetic density and are useful for maintaining a healthy body weight. The results of more than 50 intervention studies were summarized in a review evaluating the relation between energy intake, body weight, and fiber consumption. An estimated fiber intake of 14 g per day was associated with a 10% reduction of energy intake and weight loss of 2 kg in 4 months. The changes observed in body weight and energy occurred, regardless of whether the fiber source was a naturally high-fiber food or a fiber supplement.¹¹⁵ In another review of more than 60 studies, the conclusion reached was that there is solid evidence that viscous dietary fiber intake (7 g/day) helps reduce body weight and the fat mass, even in the absence of calorie restriction.¹¹⁶

Among the mechanisms by which fiber intake can aid in maintaining body weight is the gut microbiota. The gut microbiota affects the absorption of nutrients and energy homeostasis through hormones that regulate the deposit of fat into the adipocytes.¹¹⁷ Studies on animals have shown that dysbiosis of the microbiota can inhibit adenosine monophosphate kinase (AMPK), which negatively affects fatty acid oxidation, promotes lipogenesis, cholesterol and triglyceride synthesis, and the deposit of fat, producing obesity.¹¹⁸ In addition, the gut microbiota has effects on the fasting-induced adipose factor (FIAF), modulates bile acid metabolism, modulates satiety, and regulates anorexigenic hormones, such as GLP1 and PYY, through the SCFAs.¹¹⁹

Studies on humans have shown that, throughout different populations, obesity and a higher BMI, in general, are associated with a low level of bacterial diversity.^{120–122} Studies that include rural and migrant populations suggest that the transition to a low-fiber diet, derived from the westernization of

the populations, coincides with an increase in body weight, as well as a loss of gut microbiota diversity.^{123–125} The results of a longitudinal study showed that higher fiber intake was associated with greater microbiota diversity and concomitant lower weight gain in the long term.¹²⁶ Those analyses sustain the premise that fiber intake, through its effect on bacterial diversity, could help regulate body weight.

Interestingly, dietary interventions with one type of fiber, despite resulting in certain benefits on metabolic health, have been shown to not necessarily increase bacterial diversity. In contrast, in studies on humans and in vitro models, variety was found in the structures of the fiber (through the consumption of different types of plants), which was associated with greater bacterial diversity.¹²⁷ Thus, the consumption of a combination of different types and sources of fiber, as opposed to fiber intake *per se*, is posited to help increase microbial diversity, and in turn, regulate body weight.¹²⁸

Fiber promotes the growth of genera, such as lactobacilli and bifidobacteria, inducing an environment that has traditionally been referred to as "healthier". However, that is not completely clear, given that dysbiosis in obese persons has been found to be related to an increase in the phylum Firmicutes, the genus *Clostridium*, and in some species of *Lactobacillus*, signifying that not necessarily all members of the genus *Lactobacillus*, specifically in the context of obesity, have a positive connotation.¹²⁹ Albeit there is no definition of what a normal microbiota is, fiber intake has a protective effect against body weight gain and the incidence of DM that is partially mediated by the gut microbiota.¹³⁰ Strikingly, numerous studies have found great interindividual variability in the response to interventions with different kinds of fiber. Therefore, if those interventions can induce changes in the composition of the microbiota, the microbiota is thought to be able to determine how the fiber is metabolized, thus having an impact on the health of the individual.¹³¹ Studies on adults with overweight and obesity have shown that individuals with a specific gut microbiota profile can obtain greater benefit regarding body weight loss, after interventions that are rich in fiber. Individuals whose microbiota has a greater abundance of the genus *Prevotella* in relation to *Bacteroides* particularly appear to lose body weight after the intervention.^{132–134} Other genera associated with the degree of response to interventions with fiber, such as inulin, are *Akkermansia*, *Butyricoccus*, *Anaerostipes*, and *Bifidobacterium*.^{135,136} Despite the fact that the characterization of the gut microbiota is not currently carried out systematically, the above evidence could have clinical implications, in which the incorporation of gut microbiota markers could aid in improving the efficiency of nutritional therapies.¹³⁷

Dietary fiber and colon cancer

Excess protein intake leads to fermentation in the colon, with the production of compounds that have been associated with colorectal cancer, but evidence confirming that is insufficient. A simple strategy to counteract adverse effects, if there were any, would be to reduce protein intake or administer synbiotics.¹³⁸

Regardless of colorectal cancer etiology, mucosal biomarkers were reported to be reversed, with the administration of 55 g of fiber daily.¹³⁹

Epidemiologic studies provide important information on fiber intake and colorectal cancer.¹⁴⁰ In a meta-analysis of 11 prospective cohort studies, dietary fiber intake was inversely associated with the risk for both proximal and distal colon cancer.¹⁴¹ Several years earlier, a meta-analysis of 25 prospective studies had found that high total fiber intake, particularly fiber from cereals and whole grains, was associated with a lower risk for colorectal cancer.¹⁴² In a very recent meta-analysis of 22 studies, groups of adults with very high fiber intake were compared with those with very low fiber consumption. The results of the analysis suggest that dietary fiber intake could protect against rectal cancer, with a clinically relevant risk reduction.¹⁴³ Fiber not only has an effect on cancer, but on inflammatory bowel disease, as well. In non-industrialized regions of Africa, in which its inhabitants consume more than 50 g/day of fiber, the prevalence of chronic inflammatory diseases is very low.¹⁴⁰

Dietary fiber and constipation

The relation between fiber and bowel movement ease is related to certain properties of fiber, such as its capacity to retain water, increase fecal volume, increase intestinal propulsion, and reduce bowel transit time. Thus, it is important for maintaining normal bowel habit regularity. Fiber augments the food bolus, and the consequent distension of the intestine produces an increase in peristalsis.¹⁴⁴ Said increase in the bolus is the result of liquid retention between the fiber and increased bacterial density, due to fermentation. Fiber supplementation of 20–30 g/day is the usual consideration for adult patients with chronic constipation.¹⁴⁵ The use of different types of fiber for that purpose is similarly effective.¹⁴⁶ Less fermentable fibers have greater water-holding capacity and greater resistance to bacterial degradation, compared with more fermentable fibers, which is important, because bowel transit is thought to become faster, the greater the added weight. However, in an in-depth review of different interventions, bowel transit time was reduced only in individuals with initial transit times above 48 h, regardless of the type of fiber. In the same study, cereal and vegetable dietary fibers had comparable effects on fecal weight, superior to those of fruit fibers.¹⁴⁷

The beneficial effect of fiber on chronic constipation has been shown in cohort studies and in intervention studies.¹⁴⁸ In a study conducted on nurses, there was a 36% reduction in constipation in the individuals in the highest dietary fiber consumption percentile, compared with those in the lowest percentile, corresponding to a 1.8% reduction in constipation for every extra gram of fiber consumed.¹⁴⁹ The potential adverse effects of dietary fiber intake are flatulence, abdominal distension, and abdominal pain due to fiber fermentability, especially in FOS consumption, which can cause symptoms at doses as low as 10 g.¹⁵⁰ Adaptations should be made in relation to the type of fiber the patient best tolerates, and fibers with low fermentability are most likely to be better tolerated in patients with preexisting conditions associated with gas and bloating.

Dietary fiber, microbiota, and irritable bowel syndrome

The effect of fiber on the symptoms of irritable bowel syndrome (IBS) is variable and specific to the type of fiber, e.g., soluble fiber, like *Psyllium*, has shown beneficial therapeutic effects, but insoluble fiber, like wheat bran, has not. Second-line dietary therapy for IBS is a diet low in fermentable oligosaccharides, disaccharides, monosaccharides, and polyols (FODMAPs). Those fermentable carbohydrates can contribute to the increase in the production of gas and exacerbate IBS symptoms.¹⁵¹ The majority of persons do not experience important symptoms after fiber consumption, and in those that do, intolerance to FODMAPs disappears over time, as the microbiota of the host adapts to their intake.¹⁵² Nevertheless, in some patients with visceral hypersensitivity, the use of low fermentability fibers, such as methylcellulose, or those of intermediate solubility, such as *Psyllium plantago* and ispaghula, is recommended.

The fructose, lactose, fructan, sorbitol, and fructooligosaccharide FODMAPs are found in fruits, onions, garlic, legumes, and wheat. After several non-controlled studies on the effects of FODMAPs on IBS symptoms, a randomized and blinded study demonstrated the improvement in the grading of symptoms of bloating, pain, and flatulence, with the implementation of a low-FODMAP diet.¹⁵³ The same occurred in a study carried out in Mexico.¹⁵⁴ According to another analysis, the response to a low-FODMAP diet was dependent on the structure of the patient's microbiota, showing less response, the greater the dysbiosis index.¹⁵⁵

Changes in the microbiota associated with a low-FODMAP diet have been demonstrated, and so whether its long-term use is adequate is still unclear. FODMAPs should be restricted in relation to adequate symptom control.¹⁵⁶ The establishment of a low-FODMAP diet should always be carried out under the supervision of healthcare professionals trained in regard to these types of recommendations because such a diet could result in nutritional deficiencies and disorderly eating behaviors.¹⁵⁰

With respect to the pediatric population, due to the scarcity of clinical trials on the use of fiber in children with IBS, a definitive conclusion cannot be reached. Healthcare professionals should be cautious when selecting the type and dose of fiber in children and adults with IBS, to not worsen their symptoms. The use of a low-FODMAP diet is not currently recommended for the pediatric population.¹⁵⁷

The composition of the intestinal bacteria in patients with IBS differs from that of healthy subjects, with less abundance of the butyrate producers, *Erysipelotrichaceae* and *Ruminococcaceae*, than healthy children. Likewise, they have greater abundance of Methanobacteriales (methane-producing bacteria), *Lactobacillus* and *Ruminococcus*, and a decrease in *Bifidobacterium*, *Faecalibacterium*, *Erysipelotrichaceae*, and methanogens.^{158–160} When the microbiota of the patient with IBS is rebalanced due to the effect of treatment, the SCFAs it produces have a potentially beneficial effect, such as improving epithelial renewal, improving intestinal permeability, and reducing low-grade inflammation. However, more studies need to be carried out to determine the mechanisms through which fiber improves

aspects of IBS pathophysiology. There are increasingly more tests demonstrating that not only the abdominal symptoms of IBS, but also the psychiatric comorbidity that presents in a considerable number of those patients, are explained by the gut microbiota.

Dietary fiber and inflammatory bowel disease

The glycoprotein and polysaccharide-rich layer of mucus that covers the surface of the intestinal mucosa is the first line of defense between intestinal cells and the gut microbiota,¹⁶¹ and in turn, is a source of nutrition for certain gut bacteria.¹⁶² *B. thetaiotaomicron* has been shown to metabolize mucus glycans when there is a lack of dietary fiber, thinning the layer and resulting in the close contact of the bacteria with the epithelium.¹⁶³ That could explain the damage that can be caused by the deficiency of fiber in IBD and colon cancer.¹⁶⁴ SCFAs exert anti-inflammatory effects on macrophages and dendritic cells because they stimulate regulatory T cell differentiation.¹⁶⁵ Patients with IBD have lower levels of SCFAs, including butyric acid and acetic acid, compared with healthy subjects. Butyric acid could provide protection against IBD.¹⁶⁶ The scientific evidence for indicating fiber as treatment for ulcerative colitis and reservoiritis is still limited.¹⁶⁷

Dietary fiber in colostomy management

Little is known about the nutritional status and eating habits of persons with intestinal stomas and no universal dietary guidelines have been established. Many persons with stomas adjust their diet to avoid discomfort that interferes with daily life and makes them afraid to leave home, such as increased odor of gases or stool production, constipation, or leaks. Some patients avoid certain foods, especially fruits and vegetables.¹⁶⁸ The increase in fiber and liquid intake is one of the more widely used measures in patients with stomas that suffer from chronic constipation, alleviating the condition in the majority of them.¹⁶⁹ Soluble fiber supplements are frequently used if dietary measures are not sufficient, but their employment is empiric, given that there are no randomized comparative studies on the topic.¹⁷⁰ The interaction between dietary fiber and the microbiota in patients with ileostomy and colostomy has not been specifically studied.

Dietary fiber and portosystemic hepatic encephalopathy

The decrease in the consumption of foods of animal origin and the increase in vegetable proteins reduces hepatic encephalopathy (HE), albeit the mechanism is not clear. When fiber is increased in the diet, its fermentation reduces the pH of the colon, favoring the excretion of ammonia, rather than its absorption, accelerating colon transit.¹⁷¹ In cirrhosis, protein consumption should not be reduced, but rather plant-based protein, naturally associated with dietary fiber, should be administered.¹⁷² The microbiota is capable of producing the majority of neurotransmitters found in the human brain and they obviously influence neu-

rochemistry and behavior. HE is considered the prototypic brain-gut-microbiota axis disorder. Translational research indicates that certain bacteria and their manipulation can have an impact on the positive responses of brain function. The increase in fermentable fiber could reduce the absorption of ammonia in the portal system similarly to that of lactulose supplementation.¹⁷³

The use of dietary fiber as the only therapeutic measure against HE had not been studied, but a fiber-rich diet enables the concomitant increase of protein intake. A Mexican study showed that supplementation with branched-chain amino acids plus a diet with a high content of fiber and proteins is a safe intervention in patients with cirrhosis, given that it helps increase muscle mass without elevating ammonemia or fostering the development of HE.¹⁷⁴ Numerous well-designed studies have evaluated the benefit of different probiotics in the treatment of HE. Compared with placebo or no intervention, probiotics most likely improve recovery and can regulate ammonia levels in plasma, as well as quality of life, in patients with overt HE, although with no decrease in mortality.¹⁷⁵ At present, there are no studies in the literature that describe the modification of the gut microbiota in cirrhotic patients with HE, as a response to specific diets.¹⁷⁶ High quality clinical trials are needed to clarify the true potential of dietary fiber, the efficacy of probiotics, and their effect on the gut microbiota in HE.

Conclusions

Dietary fiber can induce changes in intestinal health that are directly and indirectly mediated by the gut microbiota. The use of dietary fiber by the gut microbiota depends on various factors and characteristics of the fiber, such as its fermentability, solubility, and viscosity. The type of dietary fiber influences the composition of the microbiota, given that not all bacterial species degrade all types of fiber, which can be verified by changes at the level of the phylum, family, and species. Diets low in dietary fiber can reduce the production of SCFAs, affecting their different local and systemic functions. The indirect benefits of dietary fiber impact cardiometabolic and digestive health, including some functional gastrointestinal disorders, as well as different diseases. The recommended daily intake of dietary fiber in adolescents and adults is 14 g for every 1000 kcal, in general. In pathologic cases, treatment should be individualized, with very close follow-up.

Financial disclosure

The scientific agenda, discussion, and conclusions of the present review were autonomously determined and independently drafted by the members of the group summoned by the *Asociación Mexicana de Gastroenterología (AMG)*.

For strictly logistical purposes, the AMG requested and received funding, with no conditions, from the *Instituto de Nutrición and Salud Kellogg's® (INSK)*. The authors received an honorarium for their participation.

Conflict of interest

MTAA: has received honoraria for advisory board consulting services and for conferences for Takeda®, Alfasigma®, Mayoly-Spindler®, Sanofi®, Menarini®, Carnot®, Biocodex®, BioGaia®, and Abbott® Farmacéutica.

MPMG: has received honoraria for conferences Kellogg's®, Abbott® Farmacéutica, Victus®, and Alfasigma®.

GAAA: has received funding for attending congresses and honoraria for conferences Nestlé® and Carnot®.

AMCB: has no conflict of interest.

RICS: has received honoraria for advisory board consulting services for Asofarma®, and is a speaker for Mayoly-Spindler®, Asofarma® y Chinoin®.

ACS: has received honoraria for conferences for Sanofi-Aventis®, Mead Johnson®, Takeda®, and Alexion Pharma®, and funding for clinical research studies from Alexion Pharma® and Sanofi-Aventis®.

ECA: is a member of the advisory board of Takeda® Mexico, Ferrer® Mexico, Asofarma® Mexico, Grunenthal® Mexico, and is a Speaker for Takeda® Mexico, Ferrer® Mexico, Asofarma® Mexico, Grunenthal® Mexico, Medtronic® Mexico, Asofarma® Central America and the Caribbean, Carnot® Mexico, and Siegfried Rhein® Mexico.

MFGC: has received honoraria for conferences for Grunenthal® Mexico.

VHR: has received funding to attend congresses and honoraria for conferences for Nestlé®.

MEIC: has received funding to attend congresses and has received honoraria for talks for Asofarma®, Takeda®, and Chinoin®.

JNMM: has no conflict of interest.

SMR: has received honoraria for conferences for Kellogg's® and PepsiCo®.

EOO: has received funding to attend congresses and has received honoraria for conferences and the development of materials for VitaFlo Argentina®, Nestlé®, Danone®, Abbott®, and Kellogg's®.

MRA: has received funding to attend congresses from Imed Orphan® and has received honoraria for advisory board consulting services for Abbott®.

RLRF: has received funding to attend congresses for Gilead Sciences®.

FZD: has no conflict of interest.

FZM: has received funding to attend congresses and has received honoraria for conferences and advisory board consulting services from Mead Johnson®.

RVF: has received funding to attend congresses, has received honoraria for conferences for Nestlé®, Sanofi®, Carnot®, BioGaia®, and Abbott® Farmacéutica, and has received funding for clinical research studies from Alexion Pharma® and Sanofi-Aventis®.

References

1. Holscher HD. Dietary fiber and prebiotics and the gastrointestinal microbiota. *Gut Microbes*. 2017;8:172–84, <http://dx.doi.org/10.1080/19490976.2017.1290756>.
2. Institute of Medicine (US) Panel on the definition of dietary fiber and the standing committee on the scientific evaluation of dietary reference intakes. Dietary reference intakes proposed definition of dietary fiber.

- Washington (DC): National Academies Press (US); 2001, <http://dx.doi.org/10.17226/10161>.
3. Jones JM. CODEX-aligned dietary fiber definitions help to bridge the 'fiber gap'. *Nutr J.* 2014;13:34, <http://dx.doi.org/10.1186/1475-2891-13-34>.
 4. Stephen AM, Champ MM, Cloran SJ, et al. Dietary fibre in Europe: current state of knowledge on definitions, sources, recommendations, intakes and relationships to health. *Nutr Res Rev.* 2017;30:149–90, <http://dx.doi.org/10.1017/S095442241700004X>.
 5. Korczak R, Kamil A, Fleige L, et al. Dietary fiber and digestive health in children. *Nutr Rev.* 2017;75:241–59, <http://dx.doi.org/10.1093/nutrit/nuw068>.
 6. De Filippo C, Cavalieri D, Di Paola M, et al. Impact of diet in shaping gut microbiota revealed by a comparative study in children from Europe and rural Africa. *Proc Natl Acad Sci U S A.* 2010;107:14691–6, <http://dx.doi.org/10.1073/pnas.1005963107>.
 7. Slavin J. Fiber and prebiotics: mechanisms and health benefits. *Nutrients.* 2013;5:1417–35, <http://dx.doi.org/10.3390/nu5041417>.
 8. Gray J. Dietary Fibre International Life Sciences Institute. ILSI Europe. 2006 [Accessed 19 February 2020]. Available from: <https://ils.eu/publication/dietary-fibre/>.
 9. Escudero-Álvarez E, González-Sánchez P. Fibra dietética. *Nutr Hosp.* 2006;21:61–72.
 10. Bueter CL, Specht CA, Levitz SM. Innate sensing of chitin and chitosan. *PLoS Pathog.* 2013;9:e1003080, <http://dx.doi.org/10.1371/journal.ppat.1003080>.
 11. Dahl WJ, Stewart ML. Position of the Academy of Nutrition and Dietetics: health implications of dietary fiber. *J Acad Nutr Diet.* 2015;115:1861–70, <http://dx.doi.org/10.1016/j.jand.2015.09.003>.
 12. Oliveira G, González-Molero I. An update on probiotics, prebiotics and symbiotics in clinical nutrition. *Endocrinol Nutr.* 2016;63:482–94, <http://dx.doi.org/10.1016/j.endonu.2016.07.006>.
 13. Bae SH. Diets for constipation. *Pediatr Gastroenterol Hepatol Nutr.* 2014;17:203–8, <http://dx.doi.org/10.5223/pghn.2014.17.4.203>.
 14. Anderson JW, Baird P, Davis RH, et al. Health benefits of dietary fiber. *Nutr Rev.* 2009;67:188–205, <http://dx.doi.org/10.1111/j.1753-4887.2009.00189x>.
 15. Gutiérrez JP, Rivera-Dommarco J, Shamah-Levy T, et al, Available from: <https://www.redalyc.org/articulo.oa?id=10628331033>, 2012.
 16. Shamah-Levy T, Cuevas-Nasu L, Rivera-Dommarco J, et al. Informe final de resultados. ENSANUT MC. México. 2016 [Accessed 19 February 2020]. Available from: <https://www.gob.mx/salud/documentos/encuesta-nacional-de-salud-y-nutricion-de-medio-camino-2016>
 17. Makharia G, Gibson P, Bai J, et al. Diet and the Gut. World Gastroenterology Organization Global Guidelines. 2018 [Accessed 19 February 2020]. Available from: <https://www.worldgastroenterology.org/guidelines/global-guidelines/diet-and-the-gut/diet-and-the-gut-english>
 18. Hamaker BR, Tuncil YE. A perspective on the complexity of dietary fiber structures and their potential effect on the gut microbiota. *J Mol Biol.* 2014;426:3838–50, <http://dx.doi.org/10.1016/j.jmb.2014.07.028>.
 19. Mcrorie JW, Fahey GC. A review of gastrointestinal physiology and the mechanisms underlying the health benefits of dietary fiber: Matching an effective fiber with specific patient needs. 2013;1:82–92, <http://dx.doi.org/10.5430/cns.v1n14p82>.
 20. Spiller GA. *Handbook of dietary fiber in human nutrition*. 2nd ed. Boca Raton, Florida: CRC Press, Inc.; 1993.
 21. Chaplin MF. Fiber and water binding. *Proc Nutr Soc.* 2003;62:223–7, <http://dx.doi.org/10.1079/pns2002203>.
 22. Tamargo A, Cueva C, Alvarez MD, et al. Physical effects of dietary fiber on simulated luminal flow, studied by in vitro dynamic gastrointestinal digestion and fermentation. *Food Funct.* 2019;10:3452–65, <http://dx.doi.org/10.1039/c9fo00485h>.
 23. Chen T, Chen D, Tian G, et al. Soluble fiber and insoluble fiber regulate colonic microbiota and barrier function in a piglet model. *Biomed Res Int.* 2019;2019:7809171, <http://dx.doi.org/10.1155/2019/7809171>.
 24. Jung TH, Jeon WM, Han KS. In vitro effects of dietary inulin on human fecal microbiota and butyrate production. *J Microbiol Biotechnol.* 2015;25:1555–8, <http://dx.doi.org/10.4014/jmb.1505.050778>.
 25. Vandeputte D, Falony G, Vieira-Silva S, et al. Prebiotic inulin-type fructans induce specific changes in the human gut microbiota. *Gut.* 2017;66:1968–74, <http://dx.doi.org/10.1136/gutjnl-2016-313271>.
 26. Scott KP, Martin JC, Duncan SH, et al. Prebiotic stimulation of human colonic butyrate-producing bacteria and bifidobacteria, in vitro. *FEMS Microbiol Ecol.* 2014;87:30–40, <http://dx.doi.org/10.1111/1574-6941.12186>.
 27. Valcheva R, Koleva P, Martínez I, et al. Inulin-type fructans improve active ulcerative colitis associated with microbiota changes and increased short-chain fatty acids levels. *Gut Microbes.* 2019;10:334–57, <http://dx.doi.org/10.1080/19490976.2018.1526583>.
 28. Abell GCJ, Cooke CM, Bennett CN, et al. Phylotypes related to *Ruminococcus bromii* are abundant in the large bowel of humans and increase in response to a diet high in resistant starch. *FEMS Microbiol Ecol.* 2008;66:505–15, <http://dx.doi.org/10.1111/j.1574-6941.2008.00527.x>.
 29. Ze X, Duncan SH, Louis P, et al. *Ruminococcus bromii* is a keystone species for the degradation of resistant starch in the human colon. *ISME J.* 2012;6:1535–43, <http://dx.doi.org/10.1038/ismej.2012.4>.
 30. Louis P, Flint HJ. Formation of propionate and butyrate by the human colonic microbiota. *Environ Microbiol.* 2017;19:29–41, <http://dx.doi.org/10.1111/1462-2920.13589>.
 31. Koh A, De Vadder F, Kovatcheva-Datchary P, et al. From Dietary Fiber to Host Physiology: Short-Chain Fatty Acids as Key Bacterial Metabolites. *Cell.* 2016;165:1332–45, <http://dx.doi.org/10.1016/j.cell.2016.05.041>.
 32. Morrison DJ, Preston T. Formation of short chain fatty acids by the gut microbiota and their impact on human metabolism. *Gut Microbes.* 2016;7:189–200, <http://dx.doi.org/10.1080/19490976.2015.1134082>.
 33. Baxter NT, Schmidt AW, Venkataraman A, et al. Dynamics of human gut microbiota and short-chain fatty acids in response to dietary interventions with three fermentable fibers. *mBio.* 2019;10:1–13, <http://dx.doi.org/10.1128/mBio.02566-18>.
 34. Makki K, Deehan EC, Walter J, et al. The impact of dietary fiber on gut microbiota in host health and disease. *Cell Host Microbe.* 2018;23:705–15, <http://dx.doi.org/10.1016/j.chom.2018.05.012>.
 35. Evans JM, Morris LS, Marchesi JR. The gut microbiome: the role of a virtual organ in the endocrinology of the host. *J Endocrinol.* 2013;218:R37–47, <http://dx.doi.org/10.1530/JOE-13-0131>.
 36. Bourassa MW, Alim I, Bultman SJ, et al. Butyrate, neuroepigenetics and the gut microbiome: Can a high fiber diet improve brain health? *Neurosci Lett.* 2016;625:56–63, <http://dx.doi.org/10.1016/j.neulet.2016.02.009>.
 37. Savignac HM, Corona G, Mills H, et al. Prebiotic feeding elevates central brain derived neurotrophic factor, N-methyl-D-aspartate receptor subunits

- and d-serine. *Neurochem Int.* 2013;63:756–64, <http://dx.doi.org/10.1016/j.neuint.2013.10.006>. Savignac.
38. Sudo N, Chida Y, Aiba Y, et al. Postnatal microbial colonization programs the hypothalamic-pituitary-adrenal system for stress response in mice. *J Physiol.* 2004;558:263–75, <http://dx.doi.org/10.1113/jphysiol.2004.063388>.
 39. Schmidt K, Cowen PJ, Harmer CJ, et al. Prebiotic intake reduces the waking cortisol response and alters emotional bias in healthy volunteers. *Psychopharmacology (Berl.)* 2015;232:1793–801, <http://dx.doi.org/10.1007/s00213-014-3810-0>.
 40. Yamawaki Y, Fuchikami M, Morinobu S, et al. Antidepressant-like effect of sodium butyrate (HDAC inhibitor) and its molecular mechanism of action in the rat hippocampus. *World J Biol Psych.* 2012;13:458–67, <http://dx.doi.org/10.3109/15622975.2011.585663>.
 41. Dinan TG, Stilling RM, Stanton C, et al. Collective unconscious: How gut microbes shape human behavior. *J Psychiat Res.* 2015;63:1–9, <http://dx.doi.org/10.1016/j.jpsychires.2015.02.021>.
 42. Nieto R, Kukuljan M, Silva H. BDNF and schizophrenia: from neurodevelopment to neuronal plasticity, learning, and memory. *Front Psychiatry.* 2013;4:45, <http://dx.doi.org/10.3389/fpsy.2013.00045>.
 43. Maqsood R, Stone TW. The gut-brain axis, BDNF, NMDA and CNS disorders. *Neurochem Res.* 2016;41:2819–35, <http://dx.doi.org/10.1007/s11064-016-2039-1>.
 44. Joseph J, Depp C, Shih PB, et al. Modified Mediterranean diet for enrichment of short chain fatty acids: potential adjunctive therapeutic to target immune and metabolic dysfunction in schizophrenia? *Front Neurosci.* 2017;11:155, <http://dx.doi.org/10.3389/fnins.2017.00155>.
 45. Lozupone CA, Knight R. Species divergence and the measurement of microbial diversity. *FES Microbiol Rev.* 2008;32:557–78, <http://dx.doi.org/10.1111/j.1574-6976.2008.00111.x>.
 46. Sonnenburg ED, Smits SA, Tikhonov M, et al. Diet-induced extinctions in the gut microbiota compound over generations. *Nature.* 2016;529:212–5, <http://dx.doi.org/10.1038/nature16504>.
 47. Schroeder BO, Birchenough GMH, Stahlman M, et al. Bifidobacteria or fiber protects against diet-induced microbiota-mediated colonic mucosal deterioration. *Cell Host Microbe.* 2018;23:27–40, <http://dx.doi.org/10.1016/j.chom.2017.11.004>, e7.
 48. Laitinen K, Makkala K. Overall dietary quality relates to gut microbiota diversity and abundance. *Int J Mol Sci.* 2019;20:1835, <http://dx.doi.org/10.3390/ijms20081835>.
 49. Lulli P, Mangano VD, Onori A, et al. HLA-DRB1 and -DQB1 loci in three west African ethnic groups: genetic relationship with sub-Saharan African and European populations. *Hum Immunol.* 2009;70:903–9, <http://dx.doi.org/10.1016/j.humimm.2009.07.025>.
 50. Martínez I, Stegen JC, Maldonado-Gómez MX, et al. The gut microbiota of rural papua new guineans: composition, diversity patterns, and ecological processes. *Cell Rep.* 2015;11:527–38, <http://dx.doi.org/10.1016/j.celrep.2015.03.049>.
 51. Nakayama J, Yamamoto A, Palermo-Conde LA, et al. Impact of westernized diet on gut microbiota in children on Leyte island. *Front Microbiol.* 2017;8:197, <http://dx.doi.org/10.3389/fmicb.2017.00197>.
 52. Schnorr SL, Candela M, Rampelli S, et al. Gut microbiome of the Hadza hunter-gatherers. *Nat Commun.* 2014;5:3654, <http://dx.doi.org/10.1038/ncomms4654>.
 53. Rumpagaporn P, Reuhs BL, Kaur A, et al. Structural features of soluble cereal arabinoxylan fibers associated with a slow rate of in vitro fermentation by human fecal microbiota. *Carbohydr Polym.* 2015;130:191–7, <http://dx.doi.org/10.1016/j.carbpol.2015.04.041>.
 54. Reichardt N, Duncan SH, Young P, et al. Phylogenetic distribution of three pathways for propionate production within the human gut microbiota. *ISME J.* 2014;8:1323–35, <http://dx.doi.org/10.1038/ismej.2014.14>.
 55. Hughes SA, Shewry PR, Gibson GR, et al. In vitro fermentation of oat and barley derived beta-glucans by human faecal microbiota. *FEMS Microbiol Ecol.* 2008;64:482–93, <http://dx.doi.org/10.1111/j.1574-6941.2008.00478.x>.
 56. de Vos WM, de Vos EA. Role of the intestinal microbiome in health and disease: from correlation to causation. *Nutr Rev.* 2012;70:545–56, <http://dx.doi.org/10.1111/j.1753-4887.2012.00505.x>.
 57. Allaire JM, Crowley SM, Law HT, et al. The intestinal epithelium: central coordination of mucosal immunity. *Trends Immunol.* 2018;39:677–96, <http://dx.doi.org/10.1016/j.it.2018.04.002>.
 58. Wu X, Wu Y, He L, et al. Effects of the intestinal microbial metabolite butyrate on the development of colorectal cancer. *J Cancer.* 2018;9:2510–7, <http://dx.doi.org/10.7150/jca.25324>, eCollection 2018.
 59. Walton GE, Lu C, Trogh I, et al. A randomised, double-blind, placebo controlled cross-over study to determine the gastrointestinal effects of consumption of arabinoxylan oligosaccharides enriched bread in healthy volunteers. *Nutr J.* 2012;11:36, <http://dx.doi.org/10.1186/1475-2891-11-36>.
 60. Francois IEJA, Lescroart O, Veraverbeke WS, et al. Effects of a wheat bran extract containing arabinoxylan oligosaccharides on gastrointestinal health parameters in healthy adult human volunteers: a double-blind, randomised, placebo-controlled, cross-over trial. *Br J Nutr.* 2012;108:2229–42, <http://dx.doi.org/10.1017/S0007114512000372>.
 61. Van den Abbeele P, Gérard P, Rabot S, et al. Arabinoxylans and inulin differentially modulate the mucosal and luminal gut microbiota and mucin-degradation in humanized rats. *Environ Microbiol.* 2011;13:2667–80, <http://dx.doi.org/10.1111/j.1462-2920.2011.02533.x>.
 62. Francois IEJA, Lescroart O, Veraverbeke WS, et al. Effects of wheat bran extract containing arabinoxylan oligosaccharides on gastrointestinal parameters in healthy preadolescent children. *J Pediatr Gastroenterol Nutr.* 2014;58:647–53, <http://dx.doi.org/10.1097/MPG.0000000000000285>.
 63. Snart J, Bibiloni R, Grayson T, et al. Supplementation of the diet with high-viscosity beta-glucan results in enrichment for lactobacilli in the rat caecum. *Appl Environ Microbiol.* 2006;72:1925–31, <http://dx.doi.org/10.1128/AEM.72.3.1925-1931.2006>.
 64. Jayachandran M, Chen J, Chung SSM, et al. A critical review on the impacts of beta-glucans on gut microbiota and human health. *J Nutr Biochem.* 2018;61:101–10, <http://dx.doi.org/10.1016/j.jnutbio.2018.06.010>.
 65. Vulevic J, Juric A, Walton GE, et al. Influence of galacto-oligosaccharide mixture (B-GOS) on gut microbiota, immune parameters and metabolomics in elderly persons. *Br J Nutr.* 2015;114:586–95, <http://dx.doi.org/10.1017/S0007114515001889>.
 66. Holscher HD, Doligale JL, Bauer LL, et al. Gastrointestinal tolerance and utilization of agave inulin by healthy adults. *Food Funct.* 2014;5:1142–9, <http://dx.doi.org/10.1039/c3fo60666j>.
 67. Salazar N, Dewulf EM, Neyrinck AM, et al. Inulin-type fructans modulate intestinal Bifidobacterium species populations and decrease fecal short-chain fatty acids in obese women. *Clin Nutr.* 2015;34:501–7, <http://dx.doi.org/10.1016/j.clnu.2014.06.001>.

68. Linetzky Waitzberg D, Alves Pereira CC, Logullo L, et al. Microbiota benefits after inulin and partially hydrolyzed guar gum supplementation: a randomized clinical trial in constipated women. *Nutr Hosp.* 2012;27:123–9, <http://dx.doi.org/10.1590/S0212-16112012000100014>.
69. Finegold SM, Li Z, Summanen PH, et al. Xylooligosaccharide increases bifidobacteria but not lactobacilli in human gut microbiota. *Food Funct.* 2014;5:363–445, <http://dx.doi.org/10.1039/c3fo60348b>.
70. Costabile A, Fava F, Röytiö H, et al. Impact of polydextrose on the faecal microbiota: a double-blind, crossover, placebo-controlled feeding study in healthy human subjects. *Br J Nutr.* 2012;108:471–81, <http://dx.doi.org/10.1017/S0007114511005782>.
71. Holscher HD, Bauer LL, Vishnupriya G, et al. Agave inulin supplementation affects the fecal microbiota of healthy adults participating in a randomized, double-blind, placebo-controlled, cross-over trial. *J Nutr.* 2015;145:2025–32, <http://dx.doi.org/10.3945/jn.115.217331>.
72. Dongowski G, Lorenz A, Proll J. The degree of methylation influences the degradation of pectin in the intestinal tract of rats and in vitro. *J Nutrition.* 2002;132:1935–44, <http://dx.doi.org/10.1093/jn/132.7.1935>.
73. Licht TR, Hansen M, Bergstrom A, et al. Effects of apples and specific apple components on the cecal environment of conventional rats: role of apple pectin. *BMC Microbiol.* 2010;10:13, <http://dx.doi.org/10.1186/1471-2180-10-13>.
74. Whisner CM, Martin BR, Nakatsu CH, et al. Soluble corn fiber increases calcium absorption associated with shifts in the gut microbiome: A randomized dose-response trial in free-living puberal females. *J Nutr.* 2016;146:1298–306, <http://dx.doi.org/10.3945/jn.115.227256>.
75. Le Leu RK, Winter JM, Christophersen CT, et al. Butyrylated starch intake can prevent red meat-induced O 6-methyl-2-deoxyguanosine adducts in human rectal tissue: a randomised clinical trial. *Br J Nutr.* 2015;2:220–30, <http://dx.doi.org/10.1017/S0007114515001750>.
76. Salonen A, Lahti L, Salojaarvi J, et al. Impact of diet and individual variation on intestinal microbiota composition and fermentation products in obese men. *ISME J.* 2014;8:2218–30, <http://dx.doi.org/10.1038/ismej.2014.63>.
77. Martinez I, Kim J, Duffy PR, et al. Resistant starches types 2 and 4 have differential effects on the composition of the fecal microbiota in human subjects. *PLoS One.* 2010;5:e15046, <http://dx.doi.org/10.1371/journal.pone.0015046>.
78. Ampatzoglou A, Atwal KK, Maidens CM, et al. Increased whole grain consumption does not affect blood biochemistry, body composition, or gut microbiology in healthy, low-habitual whole grain consumers. *J Nutr.* 2015;145:215–21, <http://dx.doi.org/10.3945/jn.114.202176>.
79. Freeland K, Wilson Ch, Wolever T. Adaptation of colonic fermentation and glucagon-like peptide-1 secretion with increased wheat fibre intake for 1 year in Hyperinsulinaemic human subjects. *Br J Nutr.* 2010;103:82–90, <http://dx.doi.org/10.1017/S0007114509991462>.
80. Juge N. Microbial adhesins to gastrointestinal mucus. *Trends Microbiol.* 2012;20:30–9, <http://dx.doi.org/10.1016/j.tim.2011.10.001>.
81. Forbes JD, Van Domselaar G, Bernstein CN. The gut microbiota in immune-mediated inflammatory diseases. *Front Microbiol.* 2016;7:1081, <http://dx.doi.org/10.3389/fmicb.2016.01081>.
82. Sommer F, Bäckhed F. The gut microbiota-masters of host development and physiology. *Nat Rev Microbiol.* 2013;11:227–38, <http://dx.doi.org/10.1038/nrmicro2974>.
83. Geerlings SY, Kostopoulos I, de Vos WM, et al. Akkermansia muciniphila in the human gastrointestinal tract: when, where, and how? *Microorganisms.* 2018;6:75, <http://dx.doi.org/10.3390/microorganisms6030075>.
84. Rea MC, Dobson A, O'Sullivan O, et al. Effect of broad- and narrow-spectrum antimicrobials on *Clostridium difficile* and microbial diversity in a model of the distal colon. *Proc Natl Acad Sci U S A.* 2011;108:4639–44, <http://dx.doi.org/10.1073/pnas.1001224107>.
85. Lopez-Boado YS, Wilson CL, Hooper LV, et al. Bacterial exposure induces and activates matrilysin in mucosal epithelial cells. *J Cell Biol.* 2000;148:1305–15, <http://dx.doi.org/10.1083/jcb.148.6.1305>.
86. Gevers D, Kugathasan S, Denson LA, et al. The treatment-naive microbiome in new-onset Crohn's disease. *Cell Host Microbe.* 2014;15:382–92, <http://dx.doi.org/10.1016/j.chom.2014.02.005>.
87. Chelakkot C, Ghim J, Ryu SH. Mechanisms regulating intestinal barrier integrity and its pathological implications. *Exp Mol Med.* 2018;50:103, <http://dx.doi.org/10.1038/s12276-018-0126-x>.
88. Libertucci J, Dutta U, Kaur S, et al. Inflammation-related differences in mucosa-associated microbiota and intestinal barrier function in colonic Crohn's disease. *Am J Physiol Gastrointest Liver Physiol.* 2018;315:G420–31, <http://dx.doi.org/10.1152/ajpgi.00411.2017>.
89. Rocha NP, Milagres LC, Longo GZ, et al. Association between dietary pattern and cardiometabolic risk in children and adolescents: a systematic review. *J Pediatr (Rio J).* 2017;93:214–22, <http://dx.doi.org/10.1016/j.jped.2017.01.002>.
90. Livingstone KM, McNaughton SA. Dietary patterns by reduced rank regression are associated with obesity and hypertension in Australian adults. *Br J Nutr.* 2017;117:248–59, <http://dx.doi.org/10.1017/S0007114516004505>.
91. Funtikova AN, Navarro E, Bawaked RA, et al. Impact of diet on cardiometabolic health in children and adolescents. *Nutr J.* 2015;14:118, <http://dx.doi.org/10.1186/s12937-015-0107-z>.
92. Moreno Franco B, León Latre M, Andrés Esteban EM, et al. Soluble and insoluble dietary fibre intake and risk factors for metabolic syndrome and cardiovascular disease in middle-aged adults: the AWHs cohort. *Nutr Hosp.* 2014;30:1279–88, <http://dx.doi.org/10.3305/nh.2014.30.6.7778>.
93. Liu L, Wang S, Liu J. Fiber consumption and all-cause, cardiovascular, and cancer mortalities: a systematic review and meta-analysis of cohort studies. *Mol Nutr Food Res.* 2015;59:139–46, <http://dx.doi.org/10.1002/mnfr.201400449>.
94. Hajishafiee M, Saneei P, Benisi-Kohansal S, et al. Cereal fiber intake and risk of mortality from all causes, CVD, cancer and inflammatory diseases: a systematic review and meta-analysis of prospective cohort studies. *Br J Nutr.* 2016;116:343–52, <http://dx.doi.org/10.1017/S0007114516001938>.
95. Giacco R, Costabile G, Della Pepa G, et al. A whole-grain cereal-based diet lowers postprandial plasma insulin and triglyceride levels in individuals with metabolic syndrome. *Nutr Metab Cardiovasc Dis.* 2014;24:837–44, <http://dx.doi.org/10.1016/j.numecd.2014.01.007>.
96. Gulati S, Misra A, Pandey RM. Effects of 3 g of soluble fiber from oats on lipid levels of Asian Indians - a randomized controlled, parallel arm study. *Lipids Health Dis.* 2017;16:71, <http://dx.doi.org/10.1186/s12944-017-0460-3>.
97. Kikuchi Y, Nozaki S, Makita M, et al. Effects of whole grain wheat bread on visceral fat obesity in Japanese subjects: A randomized double-blind study. *Plant Foods Hum Nutr.* 2018;73:161–5, <http://dx.doi.org/10.1007/s11130-018-0666-1>.
98. Threapleton DE, Greenwood DC, Evans CE, et al. Dietary fibre intake and risk of cardiovascular disease: Sys-

- tematic review and meta-analysis. *BMJ*. 2013;347:f6879, <http://dx.doi.org/10.1136/bmj.f6879>.
99. Ye EQ, Chacko SA, Chou EL, et al. Greater whole-grain intake is associated with lower risk of type 2 diabetes, cardiovascular disease, and weight gain. *J Nutr*. 2012;142:1304–13, <http://dx.doi.org/10.3945/jn.111.155325>.
 100. Pereira MA, O'Reilly E, Augustsson K, et al. Dietary fiber and risk of coronary heart disease: a pooled analysis of cohort studies. *Arch Intern Med*. 2004;164:370–6, <http://dx.doi.org/10.1001/archinte.164.4.370>.
 101. Cho SS, Qi L, Fahey GC Jr, et al. Consumption of cereal fiber, mixtures of whole grains and bran, and whole grains and risk reduction in type 2 diabetes, obesity, and cardiovascular disease. *Am J Clin Nutr*. 2013;98:594–619, <http://dx.doi.org/10.3945/ajcn.113.067629>.
 102. Wu K, Bowman R, Welch AA, et al. Apolipoprotein E polymorphisms, dietary fat and fiber, and serum lipids: the EPIC Norfolk study. *Eur Heart J*. 2007;28:2930–6, <http://dx.doi.org/10.1093/eurheartj/ehm482>.
 103. Bradbury KE, Crowe FL, Appleby PN, et al. Serum concentrations of cholesterol, apolipoprotein A-I and apolipoprotein B in a total of 1694 meat-eaters, fish-eaters, vegetarians and vegans. *Eur J Clin Nutr*. 2014;68:178–83, <http://dx.doi.org/10.1038/ejcn.2013.248>.
 104. Cani PD, Bibiloni R, Knauf C, et al. Changes in gut microbiota control metabolic endotoxemia-induced inflammation in high-fat diet-induced obesity and diabetes in mice. *Diabetes*. 2008;57:1470–81, <http://dx.doi.org/10.2337/db07-1403>.
 105. Yen CH, Tseng YH, Kuo YW, et al. Long-term supplementation of isomalto-oligosaccharides improved colonic microflora profile, bowel function, and blood cholesterol levels in constipated elderly people—a placebo-controlled, diet-controlled trial. *Nutrition*. 2011;27:445–50, <http://dx.doi.org/10.1016/j.nut.2010.05.012>.
 106. Nhung BT, Tuyen LD, Linh VA, et al. Rice bran extract reduces the risk of atherosclerosis in post-menopausal Vietnamese women. *J Nutr Sci Vitaminol (Tokyo)*. 2016;62:295–302, <http://dx.doi.org/10.3177/jnsv.62.295>.
 107. Jenkins DJ, Kendall CW, Nguyen TH, et al. Effect of plant sterols in combination with other cholesterol-lowering foods. *Metabolism*. 2008;57:130–9, <http://dx.doi.org/10.1016/j.metabol.2007.08.016>.
 108. Xu H, Huang X, Riserus U, et al. Dietary fiber, kidney function, inflammation, and mortality risk. *Clin J Am Soc Nephrol*. 2014;9:2104–10, <http://dx.doi.org/10.2215/CJN.02260314>.
 109. Vernay M, Aidara M, Salanave B, et al. Diet and blood pressure in 18-74-year-old adults: the French Nutrition and Health Survey (ENNS, 2006-2007). *J Hypertens*. 2012;30:1920–7, <http://dx.doi.org/10.1097/HJH.0b013e328356c59f>.
 110. Bhupathiraju SN, Tobias DK, Malik VS, et al. Glycemic index, glycemic load, and risk of type 2 diabetes: results from 3 large US cohorts and an updated meta-analysis. *Am J Clin Nutr*. 2014;100:218–32, <http://dx.doi.org/10.3945/ajcn.113.079533>.
 111. Wannamethee SG, Whincup PH, Thomas MC, et al. Associations between dietary fiber and inflammation, hepatic function, and risk of type 2 diabetes in older men: potential mechanisms for the benefits of fiber on diabetes risk. *Diabetes Care*. 2009;32:1823–5, <http://dx.doi.org/10.2337/dc09-0477>.
 112. Tripkovic L, Muirhead NC, Hart KH, et al. The effects of a diet rich in inulin or wheat fiber on markers of cardiovascular disease in overweight male subjects. *J Hum Nutr Diet*. 2015;28:476–85, <http://dx.doi.org/10.1111/jhn.12251>.
 113. Hartley L, May MD, Loveman E, et al. Dietary fiber for the primary prevention of cardiovascular disease. *Cochrane Database Syst Rev*. 2016;2016:CD011472, <http://dx.doi.org/10.1002/14651858.CD011472.pub2>.
 114. Barrett EM, Batterham MJ, Ray S, et al. Whole grain, bran and cereal fiber consumption and CVD: a systematic review. *Br J Nutr*. 2019;121:914–37, <http://dx.doi.org/10.1017/S000711451900031X>.
 115. Howarth NC, Saltzman E, Roberts SB. Dietary fiber and weight regulation. *Nutr Rev*. 2001;59:129–39, <http://dx.doi.org/10.1111/j.1753-4887.2001.tb07001.x>.
 116. Jovanovski E, Mazhar N, Komishon A, et al. Can dietary viscous fiber affect body weight independently of an energy-restrictive diet? A systematic review and meta-analysis of randomized controlled trials. *Am J Clin Nutr*. 2020;111:471–85, <http://dx.doi.org/10.1093/ajcn/nqz292>.
 117. Rosenbaum M, Knight R, Leibel RL. The gut microbiota in human energy homeostasis and obesity. *Trends Endocrinol Metab*. 2015;26:493–501, <http://dx.doi.org/10.1016/j.tem.2015.07.002>.
 118. Boulangé CL, Neves AL, Chilloux J, et al. Impact of the gut microbiota on inflammation, obesity, and metabolic disease. *Genome Med*. 2016;8:42, <http://dx.doi.org/10.1186/s13073-016-0303-2>.
 119. Kobyliak N, Virchenko O, Falalyeyeva T. Pathophysiological role of host microbiota in the development of obesity. *Nutr J*. 2016;15:43, <http://dx.doi.org/10.1186/s12937-016-0166-9>.
 120. Dahiya DK, Renuka, Puniya M, et al. Gut microbiota modulation and its relationship with obesity using prebiotic fibers and probiotics: a review. *Front Microbiol*. 2017;8:563, <http://dx.doi.org/10.3389/fmicb.2017.00563>.
 121. Moran-Ramos S, López-Contreras BE, Canizales-Quinteros S. Gut microbiota in obesity and metabolic abnormalities: A matter of composition or functionality? *Arch Med Res*. 2017;48:735–53, <http://dx.doi.org/10.1016/j.arcmed.2017.11.003>.
 122. Fan Y, Pedersen O. Gut microbiota in human metabolic health and disease. *Nat Rev Microbiol*. 2020;19:55–71, <http://dx.doi.org/10.1038/s41579-020-0433-9>.
 123. Deehan EC, Walter J. The fiber gap and the disappearing gut microbiome: implications for human nutrition. *Trends Endocrinol Metab*. 2016;27:239–42, <http://dx.doi.org/10.1016/j.tem.2016.03.001>.
 124. Vangay P, Johnson AJ, Ward TL, et al. US immigration westernizes the human gut microbiome. *Cell*. 2018;175:962–72, <http://dx.doi.org/10.1016/j.cell.2018.10.029>.
 125. Bello MGD, Knight R, Gilbert JA, et al. Preserving microbial diversity. *Science*. 2018;362:33–4, <http://dx.doi.org/10.1126/science.aau8816>.
 126. Menni C, Jackson MA, Pallister T, et al. Gut microbiome diversity and high-fibre intake are related to lower long-term weight gain. *Int J Obes (Lond)*. 2017;41:1099–105, <http://dx.doi.org/10.1038/ijo.2017.66>.
 127. McDonald D, Hyde E, Debelius JW, et al. American Gut: an open platform for citizen science microbiome research. *mSystems*. 2018;3:e00031–18, <http://dx.doi.org/10.1128/mSystems.00031-18>.
 128. Chung WSF, Walker AW, Vermeiren J, et al. Impact of carbohydrate substrate complexity on the diversity of the human colonic microbiota. *FEMS Microbiol Ecol*. 2019;95:fiy201, <http://dx.doi.org/10.1093/femsec/fiy201>.
 129. Gomes AC, Hoffmann C, Mota JF. The human gut microbiota: Metabolism and perspective in obesity. *Gut Microbes*. 2018;9:308–25, <http://dx.doi.org/10.1080/19490976.2018.1465157>.
 130. Roberfroid M, Gibson GR, Hoyles L, et al. Prebiotic effects: metabolic and health benefits. *Br J Nutr*. 2010;104:51–63, <http://dx.doi.org/10.1017/S0007114510003363>.
 131. Hughes RL, Marco ML, Hughes JP, et al. The role of the gut microbiome in predicting response to diet and the development of precision nutrition models-part i:

- overview of current methods. *Adv Nutr.* 2019;10:953–78, <http://dx.doi.org/10.1093/advances/nmz022>.
132. Hjorth MF, Roager HM, Larsen TM, et al. Pre-treatment microbial Prevotella-to-Bacteroides ratio, determines body fat loss success during a 6-month randomized controlled diet intervention. *Int J Obes (Lond).* 2018;42:580–3, <http://dx.doi.org/10.1038/ijo.2017.220>.
 133. Christensen L, Vuholm S, Roager HM, et al. Prevotella Abundance predicts weight loss success in healthy, overweight adults consuming a whole-grain diet ad libitum: a post hoc analysis of a 6-wk randomized controlled trial. *J Nutr.* 2019;149:2174–81, <http://dx.doi.org/10.1093/jn/nxz198>.
 134. Hjorth MF, Blædel T, Bendtsen LQ, et al. Prevotella-to-Bacteroides ratio predicts body weight and fat loss success on 24-week diets varying in macronutrient composition and dietary fiber: results from a post-hoc analysis. *Int J Obes (Lond).* 2019;43:149–57, <http://dx.doi.org/10.1038/s41366-018-0093-2>.
 135. de Preter V, Vanhoutte T, Huys G, et al. Baseline microbiota activity and initial bifidobacteria counts influence responses to prebiotic dosing in healthy subjects. *Aliment Pharmacol Ther.* 2008;27:504–13, <http://dx.doi.org/10.1111/j.1365-2036.2007.03588.x>.
 136. Rodríguez J, Hiel S, Neyrinck AM, et al. Discovery of the gut microbial signature driving the efficacy of prebiotic intervention in obese patients. *Gut.* 2020;69:1975–87, <http://dx.doi.org/10.1136/gutjnl-2019-319726>.
 137. Christensen L, Roager HM, Astrup A, et al. Microbial enterotypes in personalized nutrition and obesity management. *Am J Clin Nutr.* 2018;108:645–51, <http://dx.doi.org/10.1093/ajcn/nqy175>.
 138. Windey K, De Preter V, Verbeke K. Relevance of protein fermentation to gut health. *Mol Nutr Food Res.* 2012;56:184–96, <http://dx.doi.org/10.1002/mnfr.201100542>.
 139. O’Keefe SJ, Li JV, Lahti L, et al. Fat, fibre and cancer risk in African Americans and rural Africans. *Nat Commun.* 2015;6:6342, <http://dx.doi.org/10.1038/ncomms7342>.
 140. Murphy N, Norat T, Ferrari P, et al. Dietary fibre intake and risks of cancers of the colon and rectum in the European Prospective Investigation into Cancer and Nutrition (EPIC). *PLoS One.* 2012;7:e39361, <http://dx.doi.org/10.1371/journal.pone.0039361>.
 141. Ma Y, Hu M, Zhou L, et al. Dietary fiber intake and risks of proximal and distal colon cancers: a meta-analysis. *Medicine (Baltimore).* 2018;97:e11678, <http://dx.doi.org/10.1097/MD.00000000000011678>.
 142. Aune D, Chan DS, Lau R, et al. Dietary fibre, whole grains, and risk of colorectal cancer: Systematic review and dose response meta-analysis of prospective studies. *BMJ.* 2011;343:d6617, <http://dx.doi.org/10.1136/bmj.d6617>.
 143. Gianfredi V, Nucci D, Salvatori T, et al. Rectal Cancer: 20% risk reduction thanks to dietary fibre intake. Systematic review and meta-analysis. *Nutrients.* 2019;11:1579, <http://dx.doi.org/10.3390/nu11071579>.
 144. Cummings JH. 4 The Effect of Dietary Fiber on Fecal Weight and Composition. In: Spiller GA, editor. *CRC Handbook of Dietary Fiber in Human Nutrition*. Boca Raton, FL, USA: CRC Press; 1993. p. 263–333.
 145. Voderholzer WA, Schatke W, Muhlendorfer BE, et al. Clinical response to dietary fiber treatment of chronic constipation. *Am J Gastroenterol.* 1997;92:95–8.
 146. Erdogan A, Rao SSC, Thiruvaiyaru D, et al. Randomised clinical trial: mixed soluble/insoluble fibre vs. psyllium for chronic constipation. *Aliment Pharmacol Ther.* 2016;44:35–44, <http://dx.doi.org/10.1111/apt.13647>.
 147. de Vries J, Birkett A, Hulshof T, et al. Effects of cereal, fruit and vegetable fibers on human fecal weight and transit time: a comprehensive review of intervention trials. *Nutrients.* 2016;8:130, <http://dx.doi.org/10.3390/nu8030130>.
 148. Abdullah MMH, Gyles CL, Marinangeli CPF, et al. Dietary fibre intakes and reduction in functional constipation rates among Canadian adults: a cost-of-illness analysis. *Food Nutr Res.* 2015;59:28646, <http://dx.doi.org/10.3402/fnr.v59.28646>.
 149. Dukas L, Willett WC, Giovannucci EL. Association between physical activity, fiber intake, and other lifestyle variables and constipation in a study of women. *Am J Gastroenterol.* 2003;98:1790–6, <http://dx.doi.org/10.1111/j.1572-0241.2003.07591.x>.
 150. Bonnema AL, Kohlberg LW, Thomas W, et al. Gastrointestinal tolerance to chicory inulin products. *J Am Diet Assoc.* 2010;110:865–8, <http://dx.doi.org/10.1016/j.jada.2010.03.025>.
 151. Rej A, Aziz I, Tornblom H, et al. The role of diet in irritable bowel syndrome: implications for dietary advice. *J Intern Med.* 2019;286:490–502, <http://dx.doi.org/10.1111/joim.12966>.
 152. David LA, Maurice CF, Carmody RN, et al. Diet rapidly and reproducibly alters the human gut microbiome. *Nature.* 2014;505:559–63, <http://dx.doi.org/10.1038/nature12820>.
 153. Halmos EP, Power VA, Sheperd SJ, et al. A diet low in FODMAPs reduces symptoms of irritable bowel syndrome. *Gastroenterology.* 2014;146:67–75, <http://dx.doi.org/10.1053/j.gastro.2013.09.046>.
 154. Pérez-López N, Torres-López E, Zamarripa-Dorsey F. Respuesta clínica en pacientes mexicanos con síndrome de intestino irritable tratados con dieta baja en carbohidratos fermentables (FODMAP). *Rev Gastroenterol Mex.* 2015;80:180–5, <http://dx.doi.org/10.1016/j.rgm.2015.06.008>.
 155. Bennet SMP, Böhn L, Störsrud S, et al. Multivariate modelling of faecal bacterial profiles of patients with IBS predicts responsiveness to a diet low in FODMAPs. *Gut.* 2018;67:872–81, <http://dx.doi.org/10.1136/gutjnl-2016-313128>.
 156. Halmos EP, Christophersen CT, Bird AR, et al. Diets that differ in their FODMAP content alter the colonic luminal microenvironment. *Gut.* 2015;64:93–100, <http://dx.doi.org/10.1136/gutjnl-2014-307264>.
 157. Axelrod CH, Saps M. The role of in the treatment of functional gastrointestinal disorders in children nutrients. 2018;10:1650, <http://dx.doi.org/10.3390/nu10111650>.
 158. Pozuelo M, Panda S, Santiago A, et al. Reduction of butyrate- and methane-producing microorganisms in patients with Irritable Bowel Syndrome. *Sci Rep.* 2015;5:12693, <http://dx.doi.org/10.1038/srep12693>.
 159. Chong PP, Chin VK, Looi CY, et al. The microbiome and irritable bowel syndrome - a review on the pathophysiology, current research and future therapy. *Front Microbiol.* 2019;10:1136, <http://dx.doi.org/10.3389/fmicb.2019.01136>.
 160. El-Salhy M, Hatlebakk JG, Hausken T. Diet in irritable bowel syndrome (IBS): interaction with gut microbiota and gut hormones. *Nutrients.* 2019;11:1824, <http://dx.doi.org/10.3390/nu11081824>.
 161. Johansson ME, Sjövall H, Hansson GC. The gastrointestinal mucus system in health and disease. *Nat Rev Gastroenterol Hepatol.* 2013;10:352–61, <http://dx.doi.org/10.1038/nrgastro.2013.35>.
 162. Png CW, Lindén SK, Gilshenan KS, et al. Mucolytic bacteria with increased prevalence in IBD mucosa augment in vitro utilization of mucin by other bacteria. *Am J Gastroenterol.* 2010;105:2420–8, <http://dx.doi.org/10.1038/ajg.2010.281>.
 163. Sonnenburg JL, Xu J, Leip DD, et al. Glycan foraging in vivo by an intestine-adapted bacterial symbiont. *Science.* 2005;307:1955–9, <http://dx.doi.org/10.1126/science.1109051>.
 164. Desai MS, Seekatz AM, Koropatkin NM, et al. A dietary fiber-deprived gut microbiota degrades

- the colonic mucus barrier and enhances pathogen susceptibility. *Cell*. 2016;167:1339–53, <http://dx.doi.org/10.1016/j.cell.2016.10.043>.
165. Kim S, Kim JH, Park BO, et al. Perspectives on the therapeutic potential of short-chain fatty acid receptors. *BMB Rep*. 2014;47:173–8, <http://dx.doi.org/10.5483/bmbrep.2014.47.3.272>.
 166. Huda-Faujan N, Abdulmir AS, Fatimah AB, et al. The impact of the level of the intestinal short chain Fatty acids in inflammatory bowel disease patients versus healthy subjects. *Open Biochem J*. 2010;4:53–8, <http://dx.doi.org/10.2174/1874091X01004010053>.
 167. Limketkai BN, Iheozor-Ejiofor Z, Gjuladin-Hellon T, et al. Dietary interventions for induction and maintenance of remission in inflammatory bowel disease. *Cochrane Database Syst Rev*. 2019;2:CD012839, <http://dx.doi.org/10.1002/14651858.CD012839.pub2>.
 168. De Oliveira AL, Boroni Moreira AP, Pereira Netto M, et al. A cross-sectional study of nutritional status, diet, and dietary restrictions among persons with an ileostomy or colostomy. *Ostomy Wound Manage*. 2018;64:18–29.
 169. Kuczynska B, Bobkiewicz A, Studniarek A, et al. Conservative measures for managing constipation in patients living with a colostomy. *J Wound Ostomy Continence Nurs*. 2017;44:160–4, <http://dx.doi.org/10.1097/WON.0000000000000318>.
 170. Krokowicz L, Bobkiewicz A, Borejsza-Wysocki M, et al. A prospective, descriptive study to assess the effect of dietary and pharmacological strategies to manage constipation in patients with a stoma. *Ostomy Wound Manage*. 2015;61:14–22.
 171. Bosscher D, Breynaert A, Pieters L, et al. Food-based strategies to modulate the composition of the intestinal microbiota and their associated health effects. *J Physiol Pharmacol*. 2009;60:5–11.
 172. Yao CK, Fung J, Chu NHS, et al. Dietary interventions in liver cirrhosis. *J Clin Gastroenterol*. 2018;52:663–73, <http://dx.doi.org/10.1097/MCG.0000000000001071>.
 173. Amodio P, Bemeur C, Butterworth R, et al. The nutritional management of hepatic encephalopathy in patients with cirrhosis: International Society for Hepatic Encephalopathy and Nitrogen Metabolism Consensus. *Hepatology*. 2013;58:325–36, <http://dx.doi.org/10.1002/hep.26370>.
 174. Ruiz-Margáin A, Macías-Rodríguez RU, Ríos-Torres SL, et al. Effect of a high-protein, high-fiber diet plus supplementation with branched-chain amino acids on the nutritional status of patients with cirrhosis. *Rev Gastroenterol Mex*. 2018;83:9–15, <http://dx.doi.org/10.1016/j.rgmx.2017.02.005>.
 175. Dalal R, McGee RG, Riordan SM, et al. Probiotics for people with hepatic encephalopathy. *Cochrane Database Syst Rev*. 2017;2:CD008716, <http://dx.doi.org/10.1002/14651858.CD008716.pub3>.
 176. Merli M, Lebba V, Giusto M. What is new about diet in hepatic encephalopathy? *Metab Brain Dis*. 2016;31:1289–94, <http://dx.doi.org/10.1007/s11011-015-9734-5>.